


Brin - Burago - Ivanov K. Parwani.

Thm [BI]

$f: M \rightarrow \mathbb{R}^n$ p.h., $\dim(M) = 3$


\exists branching
tangent to $E^{\mathbb{R}^n} \approx E^c \oplus E^s$

Lecture 3



$$T_M = E^u \oplus E^c \oplus E^s$$

\uparrow
 ω^u



flow to separate leaves





Thm $\forall \varepsilon > 0 \exists$ foln \mathbb{F}_ε
and an onto map $h: M \rightarrow M$

- $\|h - id\| < \varepsilon$
- $\mathbb{T}\mathbb{F}_\varepsilon$ is ε -close E^{cs}
- $L \in \mathbb{F}_\varepsilon \Rightarrow h(L) \in \mathbb{F}^{cs}$

and $h|_L$ is homeo.

Reeb components




Any foln \mathcal{F} to W^u is Reebless.
In part, $\widehat{\mathcal{F}}_\varepsilon$ is Reebless.

• each leaf $L \in \widehat{\mathcal{F}}_\varepsilon$ is π_1 -injective

$$i: L \rightarrow M$$

$i_*: \pi_1(L) \rightarrow \pi_1(M)$ is an injection

- no transverse contractible cycles:



$L \in \mathcal{F}_E$ then $h(L) \in \mathcal{F}_{es}$
and the embeddings are
homotopic

$$\begin{array}{ccc} \pi_1(L) & \hookrightarrow & \pi_1(M) \\ \downarrow \cong & & \nearrow \text{injective} \\ \pi_1(h(L)) & & \end{array}$$

No transverse contractible

cycles.
Suppose $\alpha \in \pi_1 E$



$$\Delta(T\alpha, E^{\text{cs}}) > \varepsilon > 0.$$

$$\mathcal{F}_\varepsilon \quad \Delta(T\mathcal{F}_\varepsilon, E^{\text{cs}}) < \varepsilon \quad E^{\text{cs}}$$


$$\Delta(T\alpha, T\mathcal{F}_\varepsilon) > 0.$$

Prop If $f: M \rightarrow \mathbb{R}^3$ p.h. $\dim = 3$
then the universal cover
 \tilde{M} is homeo to \mathbb{R}^3 .

Proof

\tilde{M}		$\tilde{\mathbb{R}^3}$
\downarrow		
M		\mathbb{R}^3

\tilde{M}



\tilde{F}^{cs}

\tilde{L} is
simp conn



\tilde{L}

~~\mathbb{R}^2~~ or


\mathbb{R}^2

$$T\tilde{L} = E^c \oplus E^s$$

$$[\gamma] = 0 \in \pi_1(M)$$

$$\Rightarrow [\gamma] = 0 \in \pi_1(L)$$

M



Every leaf of \tilde{F}^{cs} is a plane.

\tilde{F}^{cs}

$[\tilde{\gamma}] = 0 \in \pi_1(\tilde{L})$

\tilde{F}^{cs} ←_h \tilde{F}_ε is a true foliation by planes.

(Palmeira)


$\Rightarrow \tilde{M}$ is homeo to \mathbb{R}^3 .

F^{cs} ←_h F_ε


Prop On \tilde{M} , each leaf of \tilde{F}^{cs}
intersects a leaf \tilde{W}^u at
most once.

circle $\propto \mathbb{R}E^{cs}$

proof



Define $\mathcal{U}_r(\mathcal{J}) = \{x \in \tilde{M} : \text{dist}(x, \mathcal{J}) < r\}$
 $\mathcal{J} \subset \tilde{M}$




Prop There is $C > 0$ s.t.


volume $\mathcal{U}_r(\mathcal{J}) > C \cdot \text{length } \mathcal{J}$

for any unstable curve
 $\mathcal{J} \subset \tilde{M}$.

proof Suppose J is a curve
of length $> n$



Suppose
 $x \in U_\varepsilon(J_z) \cap U_\varepsilon(J_q) \neq \emptyset$ each J_k has
 length = 1.



$U_\varepsilon(J_k) > C.$

$M = \mathbb{T}^3$ $\tilde{M} = \mathbb{R}^3$ w/ std volume.

Suppose $f: \mathbb{T}^3 \hookrightarrow \text{p.h.}$

Lift $f: \mathbb{T}^3 \hookrightarrow \text{p.h.}$ there is

a group act $f_*: \pi_1(\mathbb{T}^3) \hookrightarrow$

s.t.

$$\gamma \in \pi_1(\mathbb{T}^3) \quad \gamma: \mathbb{R}^3 \hookrightarrow \text{p.h.} \quad f_*(\gamma) \cdot \tilde{f} = \tilde{f} \circ \gamma.$$

$$f_* \hat{\pi}_* (\pi^3) \approx \mathbb{Z}^3$$

$$\exists \text{ linear map } \tilde{A}: \mathbb{R}^3 \rightarrow \mathbb{R}^3$$
$$A: \pi^3 \rightarrow \pi^3$$

$$\text{so } A_* = f_*.$$

A "linear part" of f .

A is partially hyp

A has eigenvalues

$$|\lambda_1| \leq |\lambda_2| \leq |\lambda_3|$$

want

$$|\lambda_1| < |\lambda_3|$$

and

$$|\lambda_1| < |\lambda_2| < |\lambda_3|$$

Say $|\lambda_3| \leq 1 \Rightarrow |\lambda_i| \leq 1$


$$A = \begin{pmatrix} 1 & a & c \\ 0 & 1 & b \\ 0 & 0 & 1 \end{pmatrix}$$
$$B_N = \{x \in \mathbb{R}^3 : \|x\| < N\}$$

$$\tilde{A}^n B_1 \subset B_{p(n)}$$

$$A_* = f_* \quad \|\tilde{A} - \tilde{f}\|_0 < K \text{ on } \mathbb{R}^3$$

$$\|\hat{A}^n - \hat{f}^n\| < nK$$

$$\hat{f}^n B_1 \subset B_{p(n)} + nK$$



$$a \cdot \mu^n \leq (\text{length } \tilde{f}^n) \leq \text{volume } U, (\tilde{f}^n) \\ (\mu > 1)$$

$$\leq \text{volume } B_{p(n)+nK+1} \\ \sim (p(n)+nK+1)^3$$




~~$|\lambda_3| \leq 1$~~

$$|\lambda_1| < 1 < |\lambda_3|$$

A p.h.

$$|\lambda_1| < |\lambda_2| < |\lambda_3|$$

$$|\lambda_1| < 1 < |\lambda_3|$$



If $f: \mathbb{T}^3$ is p.h.

$A: \mathbb{T}^3 \rightarrow \mathbb{T}^3$ linear and $f_* = A_*$

then A is p.h.

M non-trivial circle bundle
over \mathbb{T}^2 .

nilmanifold.

$$0 \rightarrow S^1 \rightarrow M \rightarrow \mathbb{T}^2 \rightarrow 0$$

$$0 \rightarrow \mathbb{Z} \rightarrow \pi_1(M) \rightarrow \mathbb{Z}^2 \rightarrow 0$$

$$\hat{M} = \mathcal{H}$$


Heisenberg
space.

$\hat{\pi}_1(M)$
is
nilpotent


Malcev $f: M \mathcal{S}$ nilmanifold

$$f_*: \pi_1(M) \mathcal{S}$$

\exists algebraic map $\Phi: M \mathcal{S}$
s.t. $\Phi_* = f_*$. $A \in GL(2, \mathbb{Z})$



(v, t)



$\Phi(v, t) = (Av, p(v, t))$
 p is a quadratic poly.

$$0 \rightarrow \mathbb{Z} \rightarrow \pi_1(M) \rightarrow \mathbb{Z}^2 \rightarrow 0$$

$$\downarrow \pm \text{Id}$$

$$\downarrow f_* = \Phi_*$$

$$\downarrow A$$

$$0 \rightarrow \mathbb{Z} \rightarrow \pi_1(M) \rightarrow \mathbb{Z}^2 \rightarrow 0$$

A has eigs λ and λ^{-1}

If $|\lambda| \leq 1$


In Heisenberg space

$$\text{vol}(B_R) \sim R^4$$

polynomial growth

~~$|\lambda| \leq 1$~~

$|\lambda^{-1}| < 1 < |\lambda|$



Φ is a P.h. skew product.


$A \in GL(2, \mathbb{Z})$ hyperbolic.

$$M_A = \mathbb{T}^2 \times \mathbb{R} \xrightarrow{(Ax, t) \sim (x, t+1)}$$

$$0 \rightarrow \mathbb{Z}^2 \rightarrow \pi_1(M_A) \rightarrow \mathbb{Z} \rightarrow 0$$

$\pi_1(M_A) = \mathbb{Z}^2 \rtimes_A \mathbb{Z}$

solvable but not nilpotent.



f time one map of flow

$$\text{Id} = f_* : \pi_1(M_A) \hookrightarrow$$

$$f : M_A \hookrightarrow \text{p.h.}$$

$\exists n \geq 1$ and a lift $\tilde{f}^n : \tilde{M}_A \hookrightarrow$
 so $f_*^n = \text{Id}$.