

$f: M \hookrightarrow \text{p.h.}$ M 3-mfld w solv fund gp

$$M = \mathbb{T}^3$$

f has hyp linear part

$$M = \mathbb{T}^3$$

non-hyp linear part

M is a non-trivial S^1 -bundle over \mathbb{T}^2

$M = M_A$ is a suspension manifold.

Thm If $f: \mathbb{T}^3 \rightarrow \mathbb{T}^3$ is p.h.

and its linear part $A: \mathbb{T}^3 \rightarrow \mathbb{T}^3$ is Anosov, then

f is leaf conjugate to A .

M

(f, ω_f^c)

(g, ω_g^c)

A leaf conj is a homeo

$h: M \rightarrow M$ s.t.

$L \in \omega_f^c \Rightarrow h(L) \in \omega_g^c$

and

$$h f(L) = g h(L).$$

Then If $f: \mathbb{T}^3 \rightarrow \text{p.h.}$

with hyperbolic linear
part $A: \mathbb{T}^3 \rightarrow \mathbb{T}^3$

then f is leaf conj
to A .

$$\tilde{M} = \mathbb{R}^3 \quad f: \mathbb{R}^3 \rightarrow \mathbb{R}^3$$

linear Anosov $A: \mathbb{R}^3 \rightarrow \mathbb{R}^3$


and $\text{dist.}(f, A) < \infty$

$$T\mathbb{R}^3 = \underbrace{E^u \oplus E^c}_{W^{cu}} \oplus \underbrace{E^s}_{W^{cs}}$$

linear splitting


$$T\mathbb{R}^3 = E_A^u \oplus E_A^c \oplus E_A^s$$

$W_f^{cs} \mid \exists R$ st every leaf
of W_f^{cs} is R -close to
a linear cs -plane



Same holds for W_f^{cu}
so by intersecting
leaves of W_f^c are
 \approx R -close to
linear c -lines.

GPS: $\forall x, y \in \tilde{M} \exists! p, q \in \tilde{M}$



C_f the space of center leaves on $\tilde{M} = \mathbb{R}^3$

$$C_A = \mathbb{R}^3 / E_A^c \approx \mathbb{R}^2 \quad \text{Build}$$

$C_f \approx \mathbb{R}^2$ homeo

$$H: C_f \rightarrow C_A \text{ s.t.}$$



$$Hf = AH.$$

$$CS_f \rightarrow CS_A$$

$$H^{cs}: CS_f \rightarrow CS_A$$



Guess
 $H_0: CS_f \rightarrow CS_A$



$$H_0, \quad H_n := A^{-n} H_0 f^n$$


$$H_n \rightarrow H^{cs} \quad \text{error} \rightarrow 0$$

$$\text{so } H^{cs} f = \cancel{A} H^{cs}.$$

$$CS_f \xrightarrow{H^{cs}} CS_A \quad \Bigg| \quad H: C_f \rightarrow C_A$$

$$CU_f \rightarrow CU_A \quad \Bigg| \quad Hf = AH.$$

us-pseudoleaf



$$W_f^{us}(x) = \bigcup_{y \in W_f^u(x)} W_f^s(y)$$

\mathbb{C}^0 plane.

GPS \Rightarrow intersects
each center
leaf exactly once

Want a section Σ s.t.


- intersecting
each c-leaf
once



- unit cts


- finite
dist from
 P

$\sigma \subset W_f^{ns}(x)$
compact

$z \in \mathbb{Z}^3$









~~$W_f^{us}(x) \Sigma$~~

Don't know if $W_f^{us}(x)$ is unif
cts finite dist from P.

$\mathbb{T}^3 \rightarrow \mathbb{T}^3$

Family of leaf conj
 \uparrow
 $\mathcal{H}_0 = \{h_z : z \in \mathbb{Z}^3\}$
 \downarrow
 equifcts.

h_1, h_2, h_3



Define average $\frac{1}{3}(h_1 + h_2 + h_3)$.

$H_1 = \{ \text{all averages of eqts}$
in $H_0 \}$.

equi eqts.


\downarrow
 $\text{Cube}(N) = \{ (i, j, k) : |i|, |j|, |k| < N \}$

$$h_n = \frac{1}{\#\text{Cube}(N)} \sum_{z \in \text{Cube}(N)} h_z$$

Arzela-Ascoli: $\exists h_k$

$$h_{n_k} \rightarrow h_\infty$$

h_∞ descends to a
leaf conj on π^3 .



Thm If $f: M \rightarrow S^1$ p.h.
and M is a non-triv
 S^1 -bundle over \mathbb{T}^2
then f is a skew
product topological.

\tilde{M} Heisenberg space.

$$f: \hat{M} \rightarrow \mathfrak{g}$$

$$\text{dist}_0(\hat{M}, \mathfrak{g})$$

$$\underline{\Phi}: \hat{M} \rightarrow \mathfrak{g}$$


$$\text{dist}_0(f, \underline{\Phi})$$

There are coordinates (x, y, z) for \hat{M} s.t. lie gp homo aut.



(x, y, z) for \hat{M} s.t.

$$\underline{\Phi}(x, y, z) = (\lambda'x, \lambda y, z).$$

\hat{M}



$|\Theta\rangle$



f^N

n

