

A Note on Hadwiger's Conjecture

David R. Wood*

April 25, 2013

Hadwiger's Conjecture [7] states that every K_{t+1} -minor-free graph is t -colourable. It is widely considered to be one of the most important conjectures in graph theory; see [21] for a survey. If every K_{t+1} -minor-free graph has minimum degree at most δ , then every K_{t+1} -minor-free graph is $(\delta + 1)$ -colourable by a minimum-degree-greedy algorithm. The purpose of this note is to prove a slightly better upper bound.

Lemma 1. *Fix $t \geq 2$. Assume that every K_{t+1} -minor-free graph has minimum degree at most δ , and that every K_t -minor-free graph with exactly δ vertices has an independent set of size α . Then every K_{t+1} -minor-free graph is $(\delta - \alpha + 2)$ -colourable.*

Proof. We prove, by induction on n , that every n -vertex K_{t+1} -minor-free graph is $(\delta - \alpha + 2)$ -colourable. The base case with $n = 1$ is trivial since $\delta - \alpha + 2 \geq 2$. Let G be a K_{t+1} -minor-free graph. Let v be a vertex of minimum degree d in G . Thus $d \leq \delta$. If $d = 0$ then, by induction, $G - v$ is $(\delta - \alpha + 2)$ -colourable, and by assigning to v any colour already in use, we obtain a $(\delta - \alpha + 2)$ -colouring of G , as desired. Now assume that $d \geq 1$.

Let H be the subgraph of G induced by $N(v)$. Thus H has d vertices and no K_t -minor. Let H' be the graph obtained from H by adding $\delta - d$ isolated vertices. Then H' has exactly δ vertices and H' also has no K_t -minor. By assumption, H' has an independent set of size α . Thus H has an independent set T of size $\alpha - \delta + d$.

Let G' be the graph obtained from G by contracting each edge vw where $w \in T$ into a new vertex z . Since G' is a minor of G , G' is K_{t+1} -minor-free. Since $d \geq 1$, G' has less vertices than G . By induction, G' is $(\delta - \alpha + 2)$ -colourable. Colour each vertex in T by the colour assigned to z . Colour each vertex in $G - T - v$ by the colour assigned to the same vertex in G' . Of the d neighbours of v , at least $\alpha - \delta + d$ have the same colour. Thus at most $d - (\alpha - \delta + d) + 1 = \delta - \alpha + 1$ colours are present on the neighbours of v . Hence, at least one of the $\delta - \alpha + 2$ colours is not assigned to a neighbour of v , and this colour may be assigned to v . Thus G is $(\delta - \alpha + 2)$ -colourable. \square

*School of Mathematical Sciences, Monash University, Melbourne, Australia (david.wood@monash.edu). Research supported by the Australian Research Council.

The next lemma summarises some results about independent sets in a K_{t+1} -minor-free graphs. Part (a) is the original result in this direction by Duchet and Meyniel [5]. Part (b), which is strong when t is small, is by the author [23]. Part (c), which builds upon a similar result by Fox [6], is due to Balogh and Kostochka [2]. See [3, 9, 10, 14, 15] for related results.

Lemma 2. *Every K_{t+1} -minor-free graph on n vertices has an independent set of size α , where*

- (a) $(2\alpha - 1)t \geq n$,
- (b) $(2\alpha - 1)(2t - 5) \geq 2n - 5$ for $t \geq 5$,
- (c) $(2 - \gamma)\alpha t \geq n$,

where $\gamma = (80 - \sqrt{5392})/126 \approx 0.0521 \dots$

Theorem 3. *Fix $t \geq 6$. Assume that every K_{t+1} -minor-free graph G has minimum degree at most δ . Then*

$$\chi(G) \leq \delta - \frac{2\delta - 5}{4t - 14} + \frac{3}{2}$$

and

$$\chi(G) \leq \delta \left(1 - \frac{1}{(2 - \gamma)(t - 1)} \right) + 2,$$

where $\gamma = (80 - \sqrt{5392})/126 \approx 0.0521 \dots$

Proof. Lemma 2(b) implies that every K_t -minor-free graph with δ vertices has an independent set of size α , where $(2\alpha - 1)(2(t - 1) - 5) \geq 2\delta - 5$. Thus $\alpha \geq \frac{2\delta - 5}{4t - 14} + \frac{1}{2}$. Lemma 1 implies that $\chi(G) \leq \delta - \alpha + 2 \leq \delta - \frac{2\delta - 5}{4t - 14} + \frac{3}{2}$. Similarly, Lemma 2(c) implies the second result. \square

Note that Kostochka [11, 12] and Thomason [19, 20] independently proved that $\delta \leq ct\sqrt{\log t}$ is the best possible upper bound on the minimum degree of K_{t+1} -minor-free graphs. Thus such graphs are $ct\sqrt{\log t}$ -colourable. Unfortunately, Theorem 3 makes no asymptotic improvement to this result.

We now apply these results for particular values of t .

$t = 2$: K_3 -minor-free graphs are exactly the forests, and every forest has a vertex of degree 1. Thus $\chi(G) \leq 1 - 1 + 2 = 2$ by Lemma 1, which is tight.

$t = 3$: Every K_4 -minor-free graph G has minimum degree at most 2, and every 2-vertex graph has an independent set of size 1. Thus $\chi(G) \leq 2 - 1 + 2 = 3$ by Lemma 1, which is tight.

$t = 4$: Every K_5 -minor-free graph with at least 3 vertices has at most $3n - 6$ edges [13]. Thus every K_5 -minor-free graph has average degree less than 6 and minimum degree at most 5. Every 5-vertex K_4 -minor-free graph has an independent set of size 2. Thus

$\chi(G) \leq 5 - 2 + 2 = 5$ by Lemma 1. The 4-colour theorem and Wagner's characterisation [22] implies that $\chi(G) \leq 4$.

$t = 5$: Every K_6 -minor-free graph with at least 4 vertices has at most $4n - 10$ edges [13]. Thus Every K_6 -minor-free graph has average degree less than 8, and minimum degree at most 7. Every 7-vertex K_5 -free graph has an independent set of size 2. Thus every K_6 -minor-free graph is 7-colourable by Lemma 1, which is inferior to the result by Robertson et al. [16] who proved that such graphs are 5-colourable. Note that it is open whether every K_6 -minor-free graph has minimum degree at most 6 (see [4]).

$t = 6$: Every K_7 -minor-free graph with at least 5 vertices has at most $5n - 15$ edges [13]. Thus every K_7 -minor-free graph has average degree less than 10, and minimum degree at most 9. Every 9-vertex K_6 -free graph has an independent set of size 2. Thus every K_7 -minor-free graph is 9-colourable by Lemma 1. Albar and Gonçalves [1] proved that every K_7 -minor-free graph is 8-colourable.

We conjecture that every K_7 -minor-free graph has minimum degree at most 7 (which would be tight for $K_{1,2,2,2,2}$). This conjecture would imply that every K_7 -minor-free graph is 7-colourable. Hadwiger's conjecture says that K_7 -minor-free graphs are 6-colourable.

$t = 7$: Every K_8 -minor-free graph with at least 5 vertices has at most $6n - 20$ edges [8]. Thus every K_8 -minor-free graph has average degree less than 12, and minimum degree at most 11. Every 11-vertex K_7 -free graph has an independent set of size 2. Thus every K_8 -minor-free graph is 11-colourable by Lemma 1. Albar and Gonçalves [1] proved that every K_7 -minor-free graph is 10-colourable.

Jørgensen [8] characterised the K_8 -minor-free graphs with $6n - 20$ edges as those obtained from copies of $K_{2,2,2,2,2}$ by pasting on 5-cliques. Such graphs have minimum degree 8. We conjecture that every K_8 -minor-free has minimum degree at most 8, which would imply that such graphs are 8-colourable. Hadwiger's conjecture says that K_8 -minor-free graphs are 7-colourable.

$t = 8$: Every K_9 -minor-free graph G has at most $7n - 27$ edges [17], has average degree less than 14, and minimum degree at most 13. Every 13-vertex K_8 -free graph has an independent set of size 2. Thus every K_9 -minor-free graph is 13-colourable by Lemma 1.

Song and Thomas [17] characterised the K_9 -minor-free graphs with $7n - 27$ edges as those obtained from copies of $K_{2,2,2,3,3}$ and $K_{1,2,2,2,2,2}$ by pasting on 6-cliques. Such graphs have minimum degree at most 10. We conjecture that every K_9 -minor-free has minimum degree at most 10, which would imply that such graphs are 10-colourable by Lemma 1. Hadwiger's conjecture says that K_9 -minor-free graphs are 8-colourable.

$t = 9$: Every K_{10} -minor-free graph with at least 10 vertices has at most $11n - 66$ edges [18]. Thus every K_{10} -minor-free graph has average degree less than 22, and minimum degree at most 21. By Lemma 2(b), every K_9 -minor-free 21-vertex graph has an independent set

of 3 vertices. Thus every K_{10} -minor-free graph is 20-colourable by Lemma 1 (or directly by Theorem 3).

$t = 10$: Every K_{11} -minor-free graph with at least 11 vertices has at most $13n - 89$ edges [18]. Thus every K_{11} -minor-free graph has average degree less than 26, and minimum degree at most 25. By Lemma 2(b), every K_{10} -minor-free 25-vertex graph has an independent set of 3 vertices. Thus every K_{11} -minor-free graph is 24-colourable by Lemma 1 (or directly by Theorem 3).

We emphasize that in the cases $t = 9$ and $t = 10$, while it is likely that all of the stated bounds are far from optimal¹, the utility of our approach is evident, since we may take $\alpha = 3$ in these cases.

References

- [1] BORIS ALBAR AND DANIEL GONÇALVES. On triangles in K_r -minor free graphs. 2013. arXiv: [1304.5468](https://arxiv.org/abs/1304.5468).
- [2] JÓZSEF BALOGH AND A. V. KOSTOCHKA. Large minors in graphs with given independence number. *Discrete Math.*, 311(20):2203–2215, 2011. doi: [10.1016/j.disc.2011.07.003](https://doi.org/10.1016/j.disc.2011.07.003).
- [3] JÓZSEF BALOGH, JOHN LENZ, AND HEHUI WU. Complete minors, independent sets, and chordal graphs. *Discuss. Math. Graph Theory*, 31(4):639–674, 2011. doi: [10.7151/dmgt.1571](https://doi.org/10.7151/dmgt.1571).
- [4] JÁNOS BARÁT, GWENAËL JORET, AND DAVID R. WOOD. Disproof of the list Hadwiger conjecture. *Electron. J. Combin.*, 18(1):P232, 2011. <http://www.combinatorics.org/ojs/index.php/eljc/article/view/v18i1p232>.
- [5] PIERRE DUCHET AND HENRI MEYNIEL. On Hadwiger’s number and the stability number. *Annals of Discrete Mathematics*, 13:71–73, 1982. MR: [0671905](https://dblp.org/mr/0671905).
- [6] JACOB FOX. Complete minors and independence number. *SIAM J. Discrete Math.*, 24(4):1313–1321, 2010. doi: [10.1137/090766814](https://doi.org/10.1137/090766814).
- [7] HUGO HADWIGER. Über eine Klassifikation der Streckenkomplexe. *Vierteljschr. Naturforsch. Ges. Zürich*, 88:133–142, 1943. MR: [0012237](https://dblp.org/mr/0012237).
- [8] LEIF K. JØRGENSEN. Contractions to K_8 . *J. Graph Theory*, 18(5):431–448, 1994. doi: [10.1002/jgt.3190180502](https://doi.org/10.1002/jgt.3190180502).
- [9] KEN-ICHI KAWARABAYASHI, MICHAEL D. PLUMMER, AND BJARNE TOFT. Improvements of the theorem of Duchet and Meyniel on Hadwiger’s conjecture. *J. Combin. Theory Ser. B*, 95(1):152–167, 2005. doi: [10.1016/j.jctb.2005.04.001](https://doi.org/10.1016/j.jctb.2005.04.001).

¹See Chapter 6 of the Ph.D. thesis of Song [18] for a discussion of the likely extremal graphs

- [10] KEN-ICHI KAWARABAYASHI AND ZI-XIA SONG. Independence number and clique minors. *J. Graph Theory*, 56(3):219–226, 2007. doi: [10.1002/jgt.20268](https://doi.org/10.1002/jgt.20268). MR: [2355127](https://www.ams.org/mathscinet/show-item?id=2355127).
- [11] ALEXANDR V. KOSTOCHKA. The minimum Hadwiger number for graphs with a given mean degree of vertices. *Metody Diskret. Analiz.*, 38:37–58, 1982. MR: [0713722](https://www.ams.org/mathscinet/show-item?id=0713722), Zbl: [0544.05037](https://zbmath.org/?q=sernum/0544.05037).
- [12] ALEXANDR V. KOSTOCHKA. Lower bound of the Hadwiger number of graphs by their average degree. *Combinatorica*, 4(4):307–316, 1984. doi: [10.1007/BF02579141](https://doi.org/10.1007/BF02579141). MR: [0779891](https://www.ams.org/mathscinet/show-item?id=0779891).
- [13] WOLFGANG MADER. Homomorphiesätze für Graphen. *Math. Ann.*, 178:154–168, 1968. doi: [10.1007/BF01350657](https://doi.org/10.1007/BF01350657). MR: [0229550](https://www.ams.org/mathscinet/show-item?id=0229550).
- [14] FRÉDÉRIC MAFFRAY AND HENRI MEYNIEL. On a relationship between Hadwiger and stability numbers. *Discrete Math.*, 64(1):39–42, 1987. doi: [10.1016/0012-365X\(87\)90238-X](https://doi.org/10.1016/0012-365X(87)90238-X). MR: [882610](https://www.ams.org/mathscinet/show-item?id=882610).
- [15] ANDERS SUNE PEDERSEN AND BJARNE TOFT. A basic elementary extension of the Duchet-Meyniel theorem. *Discrete Math.*, 310(3):480–488, 2010. doi: [10.1016/j.disc.2009.03.023](https://doi.org/10.1016/j.disc.2009.03.023).
- [16] NEIL ROBERTSON, PAUL D. SEYMOUR, AND ROBIN THOMAS. Hadwiger’s conjecture for K_6 -free graphs. *Combinatorica*, 13(3):279–361, 1993. doi: [10.1007/BF01202354](https://doi.org/10.1007/BF01202354). MR: [1238823](https://www.ams.org/mathscinet/show-item?id=1238823).
- [17] ZI-XIA SONG AND ROBIN THOMAS. The extremal function for K_9 minors. *J. Combin. Theory Ser. B*, 96(2):240–252, 2006. doi: [10.1016/j.jctb.2005.07.008](https://doi.org/10.1016/j.jctb.2005.07.008). MR: [2208353](https://www.ams.org/mathscinet/show-item?id=2208353).
- [18] ZIXIA SONG. *Extremal Functions for Contractions of Graphs*. Ph.D. thesis, Georgia Institute of Technology, USA, 2004. CiteSeer: [10.1.1.92.1275](https://citeseer.ist.psu.edu/viewdoc/doi/10.1.1.92.1275). MR: [2706190](https://www.ams.org/mathscinet/show-item?id=2706190).
- [19] ANDREW THOMASON. An extremal function for contractions of graphs. *Math. Proc. Cambridge Philos. Soc.*, 95(2):261–265, 1984. doi: [10.1017/S0305004100061521](https://doi.org/10.1017/S0305004100061521). MR: [0735367](https://www.ams.org/mathscinet/show-item?id=0735367), Zbl: [0551.05047](https://zbmath.org/?q=sernum/0551.05047).
- [20] ANDREW THOMASON. The extremal function for complete minors. *J. Combin. Theory Ser. B*, 81(2):318–338, 2001. doi: [10.1006/jctb.2000.2013](https://doi.org/10.1006/jctb.2000.2013). MR: [1814910](https://www.ams.org/mathscinet/show-item?id=1814910), Zbl: [1024.05083](https://zbmath.org/?q=sernum/1024.05083).
- [21] BJARNE TOFT. A survey of Hadwiger’s conjecture. *Congr. Numer.*, 115:249–283, 1996. MR: [1411244](https://www.ams.org/mathscinet/show-item?id=1411244).
- [22] KLAUS WAGNER. Über eine Eigenschaft der ebene Komplexe. *Math. Ann.*, 114:570–590, 1937. doi: [10.1007/BF01594196](https://doi.org/10.1007/BF01594196). MR: [1513158](https://www.ams.org/mathscinet/show-item?id=1513158). Zbl: [0017.19005](https://zbmath.org/?q=sernum/0017.19005).
- [23] DAVID R. WOOD. Independent sets in graphs with an excluded clique minor. *Discrete Math. Theor. Comput. Sci.*, 9(1):171–175, 2007. <http://www.dmtcs.org/dmtcs-ojs/index.php/dmtcs/article/view/617>. MR: [2335892](https://www.ams.org/mathscinet/show-item?id=2335892).