


FSSC Science Tools

Data Selection and Exploration




Data Access: File types

► *Events File (2 types)*

- *Photon files contain all needed information for science analysis*
- *Extended files contain additional information about each event that is used for specialized analysis*
 - *not necessary for most science*
- *All event classes are available in both file types*
 - *Transient - Loose quality definition, likely some background contamination*
 - *Source - Moderate quality, typically only included in transient analyses*
 - *Diffuse - Highest quality, most likely to be photons*

► *Spacecraft File*


- *Orientation and position information useful for data selection*
- *One entry every 30 seconds*



Data Access: Downloads


► Download data from:

- [*http://fermi.gsfc.nasa.gov/ssc/data/analysis/scitools/extract_latdata.html*](http://fermi.gsfc.nasa.gov/ssc/data/analysis/scitools/extract_latdata.html)
 - Allows retrieval of data for a specified region
 - Default values correspond to suggested data selections for most analysis types
- [*http://fermi.gsfc.nasa.gov/cgi-bin/ssc/LAT/WeeklyFiles.cgi*](http://fermi.gsfc.nasa.gov/cgi-bin/ssc/LAT/WeeklyFiles.cgi)
 - Weekly files contain only Diffuse-class events from the photon files
 - Weekly spacecraft files are also available
- *FTP: Can be retrieved automatically using wget*
 - Spacecraft: [*ftp://legacy.gsfc.nasa.gov/fermi/data/lat/weekly/*](ftp://legacy.gsfc.nasa.gov/fermi/data/lat/weekly/)
 - Photon: [*ftp://legacy.gsfc.nasa.gov/fermi/data/lat/allsky/*](ftp://legacy.gsfc.nasa.gov/fermi/data/lat/allsky/)




Science Analysis Tools

- ▶ *Overview of capabilities*
 - *Maximum likelihood tool—spatial-spectral analysis of region (source detection, flux)*
 - *Includes background models*
 - *Pulsars—period analysis, blind searches*
 - *Includes ephemerides DB*
 - *GRBs—temporal cuts, spectral analysis: Ftools, XSPEC*
- ▶ *Tools and documentation are released through FSSC website (<http://fermi.gsfc.nasa.gov/ssc/>)*




Science Tools: Documentation

- ▶ *Multi-tier Documentation*
 - *Full set accompanies software release*
 - *Fermi Mission Technical Handbook*
 - *Multiple levels:*
 - *Detailed analysis description ('Cicerone')*
 - *Individual tool descriptions (like fhelp)*
 - *Analysis threads (cook book examples)*





Science Tools: Structure

- ▶ “Atomic” executables
 - Allows for divergent analysis without task repetition
 - Scriptable into more complex analysis chains
- ▶ Familiar file types
 - FITS data i/o
 - IRAF style param files
 - XML source models
 - Text-based supporting files
- ▶ Familiar toolsets
 - FV, DS9, XSPEC



Science Tools: Flowchart






Parameter Files

- ▶ *Contain parameter defaults or previous values*

```
# $Header: /nfs/slac/g/glast/ground/cvs/dataSubselector/pfiles/gtselect.par,v 1.16
#
infile,f,a,"","","Input FT1 file"
outfile,f,a,"","","Output FT1 file"
ra,r,a,0,0,360,RA for new search center (degrees)
dec,r,a,0,-90,90,Dec for new search center (degrees)
rad,r,a,180,0,180, radius of new search region (degrees)
tmin,r,a,0,0,,start time (MET in s)
tmax,r,a,0,0,,end time (MET in s)
emin,r,a,30,0,,lower energy limit (MeV)
emax,r,a,300000,0,,upper energy limit (MeV)
zmax,r,a,180,0,180,maximum zenith angle value (degrees)
evclsmin,i,h,3,0,10,"Minimum event class ID"
evclsmax,i,h,3,0,10,"Maximum event class ID"
convtype,i,h,-1,-1,1,"Conversion type (-1=both, 0=Front, 1=Back)"
phasemin,r,h,0,0,1,minmum pulse phase
phasemax,r,h,1,0,1,maximum pulse phase


evtable,s,h,"EVENTS","","Event data extension"

chatter,i,h,2,0,4,Output verbosity
clobber,      b, h, yes, , , "Overwrite existing output files"
debug,        b, h, no, , , "Activate debugging mode"
gui,          b, h, no, , , "GUI mode activated"
mode,         s, h, "ql", , , "Mode of automatic parameters"
```



Science Tools: Execution

- ▶ *Parameters can be input in three ways*
 - *Command line entry - useful for scripting*
 - Allows modification of “hidden” parameters (*likely not needed for standard analyses*)
 - *Last value stored in param file for next use*
 - *Interactive prompted entry*
 - No access to hidden parameters
- ▶ *Parameter input can be mixed*
 - *%gtselect*
 - *%gtselect clobber=no*
 - *%gtselect clobber=no, infile=events.fits, outfile=events_cut.fits, etc...*



Data Selection - 1

- ▶ Event-specific cuts can be made with **gtselect**
 - Time range, energy range, position, ROI radius, zenith angle

```
[wcne-2-147-110:Meetings/Oct2009_workshop/3c454_workshop] eferrara% gtselect evclsmin=3 evclsmax=3
Input FT1 file[@0J287_indata.txt] L090923112502E0D2F37E71_PH00.fits
Output FT1 file[L090821150043E0D2F37E96_cut.fits] 3c454_ecut.fits
RA for new search center (degrees) (0:360) [133.704] 343.490616
Dec for new search center (degrees) (-90:90) [20.1085] 16.148211
radius of new search region (degrees) (0:180) [15] 15
start time (MET in s) (0:) [252460800] 266976000
end time (MET in s) (0:) [268012800] 275369897
lower energy limit (MeV) (0:) [100] 300
upper energy limit (MeV) (0:) [300000] 300000
maximum zenith angle value (degrees) (0:180) [105]
Done.
```


Hidden parameters defined
on the command line



Keyword values should
correspond EXACTLY to
values in header
(if they exist)

- ▶ Temporal cuts using spacecraft file keywords are made with **gtmktime**
 - This MUST be applied if a zenith cut was used with **gtselect**


```
[wcne-2-147-110:Meetings/Oct2009_workshop/3c454_workshop] eferrara% gtmktime
Spacecraft data file[3c454_ecut.fits] L090923112502E0D2F37E71_SC00.fits
Filter expression[DATA_QUAL==1]
Apply ROI-based zenith angle cut [yes]
Event data file[L090821150043E0D2F37E96_cut.fits] 3c454_ecut.fits
Output event file name[L090821150043E0D2F37E96_gticut.fits] 3c454_ecut_gti.fits
```



Data Selection - 2


- ▶ *Different cuts should be used for different types of data analysis*
 - *Point Source analysis*
 - *May benefit from a higher minimum energy cut due to energy-dependent PSF*
 - *Pulsar Timing analysis*
 - *Requires that spacecraft file contain more time than event file on either end*
 - *GRB analysis*
 - *Utilizes “Transient” class photons (evclsmin=1, evclsmax=3)*
- ▶ *The current set of cuts can be reviewed using **gtvcut***
- ▶ *Recommended cuts are documented at:*

http://fermi.gsfc.nasa.gov/ssc/data/analysis/documentation/Cicerone/Cicerone_Data_Exploration/Data_preparation.html



Binning for Visualization - 1

- ▶ ***gtbin*** can be used to create several useful visualization products
 - Raw counts map
 - Quick-look light curve
 - PHA1 file
- ▶ Results are in format used by other science tools like **XSPEC**
 - Includes WSC keywords for ease of viewing
- ▶ Useful to get a rough idea of the data, but do not include:
 - Exposure correction
 - Instrument responses
 - Requires Likelihood analysis for valid results




Binning for Visualization - 2

► *Making a counts map*

```
[wcne-2-147-110:Meetings/Oct2009_workshop/3c454_workshop] eferrara% gtbin  
This is gtbin version ScienceTools-v9r15p2-fssc-20090808  
Type of output file (CCUBE|CMAP|LC|PHA1|PHA2) [CCUBE] CMAP  
Event data file name[L090821150043E0D2F37E96_gticut.fits] 3c454_ecut_gti.fits  
Output file name[L090821150043E0D2F37E96_countscube.fits] 3c454_ecut_gti_cmap.fits  
Spacecraft data file name[NONE] L090923112502E0D2F37E71_SC00.fits  
Size of the X axis in pixels[120] 300 ← Here, ROI radius × image scale  
Size of the Y axis in pixels[120] 300 = size of each axis  
Image scale (in degrees/pixel)[0.25] .1  
Coordinate system (CEL - celestial, GAL -galactic) (CEL|GAL) [CEL] CEL  
First coordinate of image center in degrees (RA or galactic l)[133.704] 343.490616  
Second coordinate of image center in degrees (DEC or galactic b)[20.1085] 16.148211  
Rotation angle of image axis, in degrees[0] 0  
Projection method e.g. AIT|ARC|CAR|GLS|MER|NCP|SIN|STG|TAN:[AIT] AIT
```

Keyword values should correspond EXACTLY to values in header




Binning for Visualization - 3

► *Making a quick-look lightcurve*

```
[wcne-2-147-110:Meetings/Oct2009_workshop/3c454_workshop] eferrara% gtbin  
This is gtbin version ScienceTools-v9r15p2-fssc-20090808  
Type of output file (CCUBE|CMAP|LC|PHA1|PHA2) [CMAP] LC  
Event data file name[3c454_ecut_gti.fits]  
Output file name[3c454_ecut_gti_cmap.fits] 3c454_ecut_gti_lightcurve.fits  
Spacecraft data file name[L090923112502E0D2F37E71_SC00.fits]  
Algorithm for defining time bins (FILE|LIN|SNR) [LIN]  
Start value for first time bin in MET[0] 266976000  
Stop value for last time bin in MET[0] 275369897  
Width of linearly uniform time bins in seconds[0] 209850
```

Keyword values should correspond EXACTLY to values in header

Caution! Final bin may not be valid if full duration is not included



Binning for Visualization - 4

- ▶ *Gtbin products are easily viewable in fv or ds9*

