

# A Polynomial Bound for Untangling Geometric Planar Graphs

Prosenjit Bose · Vida Dujmović · Ferran Hurtado ·  
Stefan Langerman · Pat Morin · David R. Wood

Received: 29 November 2007 / Revised: 26 September 2008 / Accepted: 21 November 2008 /  
Published online: 12 December 2008  
© Springer Science+Business Media, LLC 2008

**Abstract** To untangle a geometric graph means to move some of the vertices so that the resulting geometric graph has no crossings. Pach and Tardos (Discrete Comput. Geom. 28(4): 585–592, 2002) asked if every  $n$ -vertex geometric planar graph can be untangled while keeping at least  $n^\epsilon$  vertices fixed. We answer this question in the affirmative with  $\epsilon = 1/4$ . The previous best known bound was  $\Omega(\sqrt{\log n / \log \log n})$ .

---

Research of P. Bose and P. Morin was partially supported by NSERC.

Research of V. Dujmović was partially supported by CRM and NSERC.

Research of F. Hurtado was supported by projects MEC MTM2006-01267 and DURSI 2005SGR00692.

Research of D.R. Wood was supported by a QEII Research Fellowship. Research initiated at the Universitat Politècnica de Catalunya, where supported by the Marie Curie Fellowship MEIF-CT-2006-023865, and by projects MEC MTM2006-01267 and DURSI 2005SGR00692.

---

P. Bose (✉) · P. Morin

School of Computer Science, Carleton University, Ottawa, Canada

e-mail: [jit@scs.carleton.ca](mailto:jit@scs.carleton.ca)

P. Morin

e-mail: [morin@scs.carleton.ca](mailto:morin@scs.carleton.ca)

V. Dujmović

Department of Mathematics and Statistics, McGill University, Montreal, Canada

e-mail: [vida@cs.mcgill.ca](mailto:vida@cs.mcgill.ca)

F. Hurtado

Departament de Matemàtica Aplicada II, Universitat Politècnica de Catalunya, Barcelona, Spain

e-mail: [Ferran.Hurtado@upc.edu](mailto:Ferran.Hurtado@upc.edu)

S. Langerman

FNRS, Département d'Informatique, Université Libre de Bruxelles, Brussels, Belgium

e-mail: [stefan.langerman@ulb.ac.be](mailto:stefan.langerman@ulb.ac.be)

D.R. Wood

Department of Mathematics and Statistics, The University of Melbourne, Melbourne, Australia

e-mail: [D.Wood@ms.unimelb.edu.au](mailto:D.Wood@ms.unimelb.edu.au)

We also consider untangling geometric trees. It is known that every  $n$ -vertex geometric tree can be untangled while keeping at least  $\Omega(\sqrt{n})$  vertices fixed, while the best upper bound was  $\mathcal{O}((n \log n)^{2/3})$ . We answer a question of Spillner and Wolff (<http://arxiv.org/abs/0709.0170>) by closing this gap for untangling trees. In particular, we show that for infinitely many values of  $n$ , there is an  $n$ -vertex geometric tree that cannot be untangled while keeping more than  $3(\sqrt{n} - 1)$  vertices fixed.

**Keywords** Geometric graphs · Untangling · Crossings

## 1 Introduction

Geometric reconfigurations consider the following fundamental problem. Given a starting and a final configuration of an object  $\mathcal{R}$ , determine if  $\mathcal{R}$  can move from the starting to the final configuration, subject to some set of movement rules. An object can be a set of disks in the plane, or a graph representing a protein, or a robot's arm, for example. Typical movement rules include maintaining connectivity of the object and avoiding collisions or crossings.

In this paper we study the problem where the object is a planar graph<sup>1</sup>  $G$ . The starting configuration is a drawing of  $G$  in the plane with vertices as distinct points and edges as straight-line segments (and possibly many crossings). Our goal is to relocate as few vertices of  $G$  as possible in order to remove all the crossings, that is, to reconfigure  $G$  to some straight line crossing-free drawing of  $G$ . More formally, a *geometric graph* is a graph whose vertices are distinct points in the plane (not necessarily in general position) and whose edges are straight-line segments between pairs of points. If the underlying combinatorial graph of  $G$  belongs to a class of graphs  $\mathcal{K}$ , then we say that  $G$  is a *geometric  $\mathcal{K}$  graph*. For example, if  $\mathcal{K}$  is the class of planar graphs, then  $G$  is a geometric planar graph. Where it causes no confusion, we do not distinguish between the geometric graph and its underlying combinatorial graph. Two edges in a geometric graph *cross* if they intersect at some point other than a common endpoint. A geometric graph with no pair of crossing edges is called *crossing-free*.

Consider a geometric graph  $G$  with vertex set  $V(G) = \{p_1, \dots, p_n\}$ . A crossing-free geometric graph  $H$  with vertex set  $V(H) = \{q_1, \dots, q_n\}$  is called an *untangling* of  $G$  if for all  $i, j \in \{1, 2, \dots, n\}$ ,  $q_i$  is adjacent to  $q_j$  in  $H$  if and only if  $p_i$  is adjacent to  $p_j$  in  $G$ . Furthermore, if  $p_i = q_i$  then we say that  $p_i$  is *fixed*, otherwise we say that  $p_i$  is *free*. If  $H$  is an untangling of  $G$  with  $k$  vertices fixed, then we say that  $G$  can be *untangled* while keeping  $k$  vertices fixed. Clearly only geometric planar graphs can be untangled. Moreover, since every planar graph is isomorphic to some crossing-free geometric graph [6, 16], trivially every geometric planar graph can be untangled while keeping at least 2 vertices fixed. For a geometric graph  $G$ , let  $\text{fix}(G)$  denote the maximum number of vertices that can be fixed in an untangling of  $G$ .

<sup>1</sup>We consider graphs that are simple, finite, and undirected. The vertex set of a graph  $G$  is denoted by  $V(G)$ , and its edge set by  $E(G)$ . The subgraph of  $G$  induced by a set of vertices  $S \subseteq V(G)$  is denoted by  $G[S]$ .  $G \setminus S$  denotes  $G[V(G) \setminus S]$ .

At the 5th Czech–Slovak Symposium on Combinatorics in Prague in 1998, Mamoru Watanabe asked if every geometric cycle (that is, all polygons) can be untangled while keeping at least  $\varepsilon n$  vertices fixed, for some  $\varepsilon > 0$ . Pach and Tardos [11] answered that question in the negative by providing an  $\mathcal{O}((n \log n)^{2/3})$  upper bound on the number of fixed vertices. Furthermore, they proved that every geometric cycle can be untangled while keeping at least  $\sqrt{n}$  vertices fixed. This lower bound has recently been improved to  $\Omega(n^{2/3})$  by Cibulka [2].

Pach and Tardos [11] asked if every geometric planar graph can be untangled while keeping  $n^\varepsilon$  vertices fixed, for some  $\varepsilon > 0$ . In recent work, Spillner and Wolff [14] showed that geometric planar graphs can be untangled while keeping  $\Omega(\sqrt{\log n / \log \log n})$  vertices fixed. The best known bound before that was 3 [7, 15]. In Sect. 4, we answer the question of Pach and Tardos [11] in the affirmative and provide the first polynomial lower bound for untangling geometric planar graphs. Specifically, our main result is that every  $n$ -vertex geometric planar graph can be untangled while keeping  $(n/3)^{1/4}$  vertices fixed.

There has also been considerable interest in untangling specific classes of geometric planar graphs. Spillner and Wolff [14] studied the untangling of geometric outerplanar graphs and showed that they can be untangled while keeping  $\Omega(\sqrt{n})$  vertices fixed; and that for every sufficiently large  $n$ , there is an  $n$ -vertex outerplanar graph that cannot be untangled while keeping more than  $2\sqrt{n-1} - 1$  vertices fixed. Thus  $\Theta(\sqrt{n})$  is the tight bound for outerplanar graphs. A  $\sqrt{n/3}$  lower bound for trees was shown by Goac et al. [7]. The best known upper bound for trees was  $\mathcal{O}((n \log n)^{2/3})$ , which was in fact proved for geometric paths, by Pach and Tardos [11].<sup>2</sup> We answer a question posed by Spillner and Wolff [14] and close the gap for trees by showing that for infinitely many values of  $n$ , there is a forest of stars that cannot be untangled while keeping more than  $3(\sqrt{n} - 1)$  vertices fixed. This result is proved in Sect. 5. In addition, in Sect. 3, we prove that every geometric tree can be untangled while keeping  $\sqrt{n/2}$  vertices fixed. Note that the same result was independently obtained by Goac et al. [7]. We conclude the paper with some open problems.

Note that our definition of a geometric graph allows for collinear vertices. The same definition is used by Goac et al. [7] and Spillner and Wolff [14], but not by Pach and Tardos [11]. While allowing collinearities makes our lower bounds stronger, it makes our upper bound (for trees) weaker. In particular, the geometric trees, for which we prove a  $3(\sqrt{n} - 1)$  upper bound, have all their vertices on a line. We conjecture a  $\text{fix}(T) \leq \mathcal{O}(\sqrt{n})$  bound for certain geometric trees  $T$  with  $V(T)$  in general position. Very recently, Kang et al. [9] proved a bound of  $\mathcal{O}(\sqrt{n}2^{\alpha(n)})$  in this setting, where  $\alpha(\cdot)$  is the inverse Ackermann function.

Untangling graphs has also been the topic of [9, 10, 13, 15]. Goac et al. [7] also studied the computational complexity of the related optimization problems and showed various hardness results.

<sup>2</sup>Pach and Tardos [11] actually proved this upper bound for geometric cycles, but their method readily applies for geometric paths.

## 2 Lower Bounds: A Useful Lemma

When proving lower bounds, our goal will be to show that given any geometric planar graph  $G$ , we can find a large subset  $R$  of vertices of  $G$  such that  $G$  can be untangled while keeping  $R$  fixed. The following geometric lemma simplifies this task by allowing us to concentrate on the case in which each vertex in  $R$  is on the  $y$ -axis. This lemma will be useful both for untangling geometric trees in Sect. 3 and for untangling general geometric planar graphs in Sect. 4.

**Lemma 1** *Let  $\overline{G}$  be an untangling of some geometric planar graph  $G$ . Let  $R$  be a set of vertices of  $G$  such that each vertex of  $R$  is on the  $y$ -axis in  $\overline{G}$  and has the same  $y$ -coordinate in  $\overline{G}$  as in  $G$ . Then there exists an untangling  $\overline{G'}$  of  $G$  in which the vertices in  $R$  are fixed.*

*Proof* The proof uses the fact that it is possible to perturb the vertices of a crossing-free geometric graph without introducing crossings. More precisely, for any crossing-free geometric graph, there exists a value  $\varepsilon > 0$  such that each vertex can be moved a distance of at most  $\varepsilon$ , and the resulting geometric graph is also crossing-free. The maximum value  $\varepsilon$  for which this property holds is called the *tolerance* of the arrangement of segments. This concept, both for the geometric realization and the combinatorial meaning of the graphs, was systematically studied in [1, 12].

Consider the untangling  $\overline{G}$  of  $G$  and let  $\varepsilon > 0$  be the tolerance of  $\overline{G}$ . Let  $X$  denote the maximum absolute value of an  $x$ -coordinate in  $G$  of a vertex in  $R$ . Let  $\overline{G''}$  be the geometric graph obtained from  $\overline{G}$  as follows. For each vertex  $v \in R$  positioned at  $(x, y)$  in  $G$ , move  $v$  from  $(0, y)$  in  $\overline{G}$  to  $(x\varepsilon/X, y)$  in  $\overline{G''}$ . The vertices not in  $R$  are unmoved. So each vertex moves a distance of at most  $\varepsilon$ , and  $\overline{G''}$  is crossing-free. Scale  $\overline{G''}$  by multiplying the  $x$ -coordinates of all vertices in  $\overline{G''}$  by  $X/\varepsilon$  to obtain a crossing-free geometric graph  $\overline{G'}$ . Then every vertex of  $R$  has the same location in  $\overline{G'}$  as it does in  $G$ . Thus  $\overline{G'}$  is an untangling of  $G$  that keeps the vertices of  $R$  fixed.  $\square$

## 3 Trees: Lower Bound

In this section we prove a lower bound for untangling geometric trees. The proof also provides a warm up to our main result, the polynomial lower bound for planar graphs.

**Theorem 1** *Every  $n$ -vertex geometric tree  $T$  can be untangled while keeping at least  $\sqrt{n/2}$  vertices fixed. That is,  $\text{fix}(T) \geq \sqrt{n/2}$ .*

In a vertex 2-coloring of  $T$ , the largest of the two color classes has at least  $n/2$  vertices. Therefore, the following lemma, coupled with Lemma 1, implies Theorem 1.

**Lemma 2** *Let  $T$  be an  $n$ -vertex geometric tree whose vertices are 2-colored. Let  $S$  be one of the two color classes. Then there exists a set  $R$  of vertices in  $T$  such that  $|R| \geq \sqrt{|S|}$ , and there is an untangling  $\overline{T}$  of  $T$  in which each vertex in  $R$  is on the  $y$ -axis and has the same  $y$ -coordinate in  $\overline{T}$  as in  $T$ .*

*Proof* Root  $T$  at any vertex and order its vertices  $\sigma := (v_1, \dots, v_n)$  based on a post-order traversal of  $T$ .

While we make no general position assumption on the vertices of  $T$ , we may assume, by a suitable rotation, that no pair of vertices of  $T$  have the same  $y$ -coordinate. Let  $R$  be a largest ordered subset  $R \subseteq S$  such that the  $y$ -coordinates of the vertices of  $R$  are either monotonically increasing or monotonically decreasing in  $\sigma$ . By the Erdős–Szekeres Theorem [5],  $|R| \geq \sqrt{|S|}$ . Without loss of generality, assume that  $R$  is monotonically increasing.

Let  $T'$  be a geometric tree obtained from  $T$  as follows. For each vertex  $v \in R$  positioned at  $(x, y)$  in  $T$ , move  $v$  to  $(0, y)$  in  $T'$ . Move each vertex in  $S \setminus R$  from its position in  $T$  to the  $y$ -axis, such that all the vertices in  $S$  appear in  $T'$  in the order  $\sigma$  on the  $y$ -axis. The vertices in  $V(T) \setminus S$  remain fixed. To complete the proof of the lemma, it remains to show how to untangle  $T'$  while keeping  $S$  fixed. We prove that by induction on  $i$ , with the following induction hypothesis.

Let  $T_i := T'[\{v_1, \dots, v_i\}]$ . For each  $i \in \{1, \dots, n\}$ , there is an untangling  $\overline{T}_i$  of  $T_i$  such that  $S \cap V(T_i)$  is fixed, and


(1) for all  $j \in \{2, \dots, i\}$ , the  $y$ -coordinate of  $v_j$  is greater than the  $y$ -coordinate of  $v_{j-1}$ .

It remains to show how to untangle  $T'$  while keeping  $S$  fixed, as illustrated in Fig. 1.

We first prove that such an untangling has the following useful property, where the *right ray* at  $(x, y)$  is the open half-line containing all the points  $\{(x', y) : x' > x\}$ .

(2) For all  $j \in \{1, \dots, i\}$  such that the parent,  $v_p$ , of  $v_j$  is not in  $T_i$ , the right ray at  $v_j$  does not intersect  $\overline{T}_i$ .

Let  $y_j$  denote the  $y$ -coordinate of  $v_j$  in  $\overline{T}_i$ . Let  $P$  be the path from  $v_p$  to the root,  $v_n$ . Suppose that some edge  $e$  of  $T_i$  crosses the right ray at  $v_j$  in  $\overline{T}_i$ . Thus  $y_j$



**Fig. 1** An untangling of the complete binary tree of depth 4. Vertices in  $S$  are depicted by squares

is between the  $y$ -coordinates of the endpoints of  $e$ . By condition (1), the index in  $\sigma$  of one endpoint of  $e$  is less than  $j$ , and the index in  $\sigma$  of the other endpoint of  $e$  is greater than  $j$ . Since  $\sigma$  is a post-order numbering,  $e$  has one endpoint in  $P$ . Since  $v_p$  is not in  $T_i$ , the remaining vertices of  $P$  are also not in  $T_i$ . Thus no such edge  $e$  is in  $T_i$ . This proves that (1) implies (2).

We are now ready to prove the lemma by induction. For  $i = 1$ , the statement is true trivially. Assume now that  $i > 1$  and that the statement is true for  $i - 1$ . There are two cases to consider:  $v_i \in S$  and  $v_i \notin S$ .

Consider first the case that  $v_i \notin S$ . Since  $S$  is a color class in a 2-coloring of  $T$ , each child of  $v_i$ , if any, is in  $S$  and thus is on the  $y$ -axis. Assign a  $y$ -coordinate to  $v_i$  that is greater than the  $y$ -coordinate of each vertex in  $\overline{T_{i-1}}$  and less than the  $y$ -coordinate of each vertex in  $S \setminus V(T_{i-1})$ . This ensures that condition (1) is maintained in  $\overline{T_i}$ . Assign a positive  $x$ -coordinate to  $v_i$  such that  $\overline{T_i}$  is crossing-free. Condition (2) on  $T_{i-1}$  guarantees that this is always possible.


Now assume the case that  $v_i \in S$ . We start with an observation. Consider a vertex  $v \in V(\overline{T_{i-1}}) \setminus S$  whose parent is not in  $\overline{T_{i-1}}$ . Let the coordinates of  $v$  in  $\overline{T_{i-1}}$  be  $(x, y)$ . Each child of  $v$  is in  $S$  and thus lies on the  $y$ -axis. Denote their  $y$ -coordinates by  $y_1, \dots, y_d$ . By condition (2), for each  $i \in \{1, \dots, d\}$ , the right ray at  $(0, y_i)$  can only be intersected by an edge incident to  $v$  in  $\overline{T_{i-1}}$ . Thus  $v$  can be moved to any position  $(x', y)$ ,  $x' \geq 0$ , and the resulting untangling of  $T_{i-1}$  still satisfies the two conditions. We are now ready to untangle  $\overline{T_i}$ . Vertex  $v_i$  is fixed, and thus its position in  $\overline{T_i}$  is predetermined. None of its children are in  $S$ . Thus we are allowed to move any child of  $v_i$  from its position in  $\overline{T_{i-1}}$  to a new position. By the above observation it is possible to move each child  $w$  of  $v_i$  (one by one, in the decreasing order of their  $y$ -coordinates) so that the resulting untangling  $\overline{T'_{i-1}}$  of  $T_{i-1}$  satisfies conditions (1) and (2), and so that the open segment  $\overline{wv_i}$  does not intersect  $\overline{T'_{i-1}}$ . Connect  $v_i$  by a segment to each of its children in  $\overline{T'_{i-1}}$ . Then the resulting untangling  $\overline{T_i}$  is crossing-free. Condition (1) is maintained since all the vertices of  $T_{i-1}$  have smaller  $y$ -coordinate than  $v_i$  in  $\overline{T_i}$ . □

### 4 Planar Graphs: Lower Bound

Let  $G$  be an  $n$ -vertex geometric planar graph. In this section we prove that  $G$  can be untangled while keeping  $(n/3)^{1/4}$  vertices fixed (as stated in Theorem 2 below). It suffices to prove this theorem for edge-maximal geometric planar graphs. Thus, for the remainder of this section, assume that  $G$  is edge-maximal.<sup>3</sup>

Let  $\mathcal{E}$  be an embedded planar graph isomorphic to  $G$ . Each face of  $\mathcal{E}$  is bounded by a 3-cycle. Canonical orderings of embedded edge-maximal planar graphs were introduced by de Fraysseix et al. [3]. They proved that  $\mathcal{E}$  has a vertex ordering  $\sigma = (v_1 := x, v_2 := y, v_3, \dots, v_n := z)$ , called a *canonical ordering*, with the following properties. Define  $G_i$  to be the embedded subgraph of  $\mathcal{E}$  induced by  $\{v_1, v_2, \dots, v_i\}$ .

<sup>3</sup>A planar graph  $H$  is edge-maximal (also called, a *triangulation*) if for all  $vw \notin E(H)$ , the graph resulting from adding  $vw$  to  $H$  is not planar.



**Fig. 2** **a** Canonical ordering of  $\mathcal{E}$ . **b** Frame  $\mathcal{F}$  of  $\mathcal{E}$ . Vertices forming a largest antichain in  $\langle \mathcal{F} \rangle$ , that is, the vertices in  $S$ , are depicted by *squares*

Let  $C_i$  be the subgraph of  $\mathcal{E}$  induced by the edges on the boundary of the outer face of  $G_i$ . Then

- $x, y$ , and  $z$  are the vertices on the outer face of  $\mathcal{E}$ .
- For each  $i \in \{3, 4, \dots, n\}$ ,  $C_i$  is a cycle containing  $xy$ .
- For each  $i \in \{3, 4, \dots, n\}$ ,  $G_i$  is biconnected and *internally 3-connected*; that is, removing any two interior vertices of  $G_i$  does not disconnect it.
- For each  $i \in \{3, 4, \dots, n\}$ ,  $v_i$  is a vertex of  $C_i$  with at least two neighbors in  $C_{i-1}$ , and these neighbors are consecutive on  $C_{i-1}$ .

For example, the ordering in Fig. 2a is a canonical ordering of the depicted embedded graph  $\mathcal{E}$ .

We now introduce a new combinatorial structure that is critical to the proof of Theorem 2. A *frame*  $\mathcal{F}$  of  $\mathcal{E}$  is the oriented subgraph of  $\mathcal{E}$  with vertex set  $V(\mathcal{F}) := V(\mathcal{E})$ , where:

- $xy$  is in  $E(\mathcal{F})$  and is oriented from  $x$  to  $y$ .
- For each  $i \in \{3, 4, \dots, n\}$  in the canonical ordering  $\sigma$  of  $\mathcal{E}$ , edges  $pv_i$  and  $v_i p'$  are in  $E(\mathcal{F})$ , where  $p$  and  $p'$  are the first and the last neighbors, respectively, of  $v_i$  along the path in  $C_{i-1}$  from  $x$  to  $y$  not containing edge  $xy$ . Edge  $pv_i$  is oriented from  $p$  to  $v_i$ , and edge  $v_i p'$  is oriented from  $v_i$  to  $p'$ , as illustrated in Fig. 2b. We call  $p$  the *left predecessor* of  $v$  and  $p'$  the *right predecessor* of  $v$ .

We also say that  $\mathcal{F}$  is a frame of  $G$ . By definition,  $\mathcal{F}$  is a directed acyclic graph with one source  $x$ , and one sink  $y$ .  $\mathcal{F}$  defines a partial order  $\langle \mathcal{F} \rangle$  on  $V(\mathcal{F})$ , where  $v \langle \mathcal{F} \rangle w$  whenever there is a directed path from  $v$  to  $w$  in  $\mathcal{F}$ .

The remainder of this section is dedicated to proving the following two lemmas, which readily imply the desired result, as shown in the proof of Theorem 2 below.

**Lemma 3** Every  $n$ -vertex geometric planar graph  $G$  whose partial order  $<_{\mathcal{F}}$  associated with a frame  $\mathcal{F}$  of  $G$  has a chain<sup>4</sup> of size  $\ell$  can be untangled while keeping  $\sqrt{\ell/3}$  vertices fixed.

**Lemma 4** Every  $n$ -vertex geometric planar graph  $G$  whose partial order  $<_{\mathcal{F}}$  associated with a frame  $\mathcal{F}$  of  $G$  has an antichain of size  $t$  can be untangled while keeping  $\sqrt{t}$  vertices fixed.

**Theorem 2** Every  $n$ -vertex geometric planar graph  $G$  can be untangled while keeping at least  $(n/3)^{1/4}$  vertices fixed. That is,  $\text{fix}(G) \geq (n/3)^{1/4}$ .

*Proof* Let  $\mathcal{F}$  be a frame of  $G$ , and let  $<_{\mathcal{F}}$  be its associated partial order. If  $<_{\mathcal{F}}$  has a chain of size at least  $\sqrt{3n}$ , then we are done by Lemma 3. Otherwise, by Dilworth's theorem [4],  $<_{\mathcal{F}}$  has a partition into at most  $\sqrt{3n}$  antichains. By the pigeon-hole principle there is an antichain in that partition that has at least  $\frac{n}{\sqrt{3n}}$  vertices, which completes the proof, by Lemma 4.  $\square$

#### 4.1 Big Chain: Proof of Lemma 3

A *chord* of a cycle  $C$  is an edge that has both endpoints in  $C$  but itself is not an edge of  $C$ . Consider a cycle  $C$  in an embedded planar graph  $\mathcal{E}$ .  $C$  is called *externally chordless* if each chord of  $C$  is embedded inside of  $C$  in  $\mathcal{E}$ . The following theorem is by Spillner and Wolff [14], although they state it in a slightly different form; see Theorem 2 in [14].

**Theorem 3** [14] Let  $G$  be a geometric planar graph and  $\mathcal{E}$  an embedded planar graph isomorphic to  $G$ . If  $\mathcal{E}$  has an externally chordless cycle on  $\ell$  vertices, then  $G$  can be untangled while keeping at least  $\sqrt{\ell/3}$  vertices fixed.

**Lemma 5** Consider any directed path on at least three vertices from  $x$  to  $y$  in  $\mathcal{F}$ . The cycle comprised of that path and edge  $xy$  is externally chordless in  $\mathcal{E}$ .

*Proof* Denote the cycle in question by  $C$ , and denote the directed path between  $x$  and  $y$  in  $C$  not containing edge  $xy$  by  $P$ . Consider a chord  $v_i v_j$  of  $C$ . Without loss of generality,  $v_i <_{\sigma} v_j$  in the canonical ordering  $\sigma$ . Thus  $v_i$  is in  $G_{j-1}$ , and  $v_i v_j$  is an edge of  $G_j$ . The neighbors of  $v_j$  in  $G_{j-1}$  appear consecutively along the boundary  $C_{j-1}$  of  $G_{j-1}$ . Let  $x_1, \dots, x_d$  be the neighbors of  $v_j$  in left-to-right order on  $C_{j-1}$ . Thus  $x_1 v_j$  and  $v_j x_d$  are arcs in  $\mathcal{F}$ . Let  $u v_j$  and  $v_j w$  be the incoming and outgoing arcs in  $P$  at  $v_j$ . Then the counterclockwise order of edges incident to  $v_j$  in  $\mathcal{E}$  is  $(u, \dots, x_1, \dots, x_d, \dots, w, \dots)$ . In particular, each edge  $v_j x_{\ell}$  is contained in the closure of the interior of  $C$ . Now  $v_i = x_{\ell}$  for some  $\ell \in [1, d]$ . Thus  $v_i v_j$  is an internal chord of  $C$ .  $\square$

<sup>4</sup>Recall that a *chain* (antichain) in a partial order is a subset of its elements that are pairwise comparable (incomparable).

This lemma, coupled with Theorem 3, implies Lemma 3, as demonstrated below.

*Proof of Lemma 3* If  $\ell < 3$ , the claim follows trivially. Assume now that  $\ell \geq 3$ . Since  $\prec_{\mathcal{F}}$  has a chain of size  $\ell$ ,  $\prec_{\mathcal{F}}$  has a maximal chain of size  $\ell' \geq \ell$ . Every maximal chain in  $\prec_{\mathcal{F}}$  is a path from  $x$  to  $y$  in  $\mathcal{F}$ . Therefore, Lemma 5 implies that  $\mathcal{E}$  contains an externally chordless cycle on  $\ell'$  vertices, and the result follows from Theorem 3.  $\square$

### 4.2 Big Antichain: Proof of Lemma 4

For each vertex  $v \in V(\mathcal{F})$ , we define  $\text{Lroof}(v)$  and  $\text{Rroof}(v)$  as the following directed paths in  $\mathcal{F}$

$$\begin{aligned} \text{Lroof}(v_1) &:= \emptyset & \text{and} & & \text{Rroof}(v_1) &:= \emptyset, \\ \text{Lroof}(v_2) &:= \emptyset & \text{and} & & \text{Rroof}(v_2) &:= \emptyset. \end{aligned}$$

For each  $i \in \{3, \dots, n\}$ , define  $\text{Lroof}(v_i)$  and  $\text{Rroof}(v_i)$  recursively, as follows

$$\begin{aligned} \text{Lroof}(v_i) &:= \text{Lroof}(p) \cup \{pv_i\}, & \text{and} \\ \text{Rroof}(v_i) &:= \{v_i p'\} \cup \text{Rroof}(p'), \end{aligned}$$

where  $p$  is the left and  $p'$  the right predecessor of  $v_i$ . Finally, define the *roof* of  $v_i$  to be  $\text{roof}(v_i) := \text{Lroof}(v_i) \cup \text{Rroof}(v_i)$ .

Note that for each  $i \in \{3, \dots, n\}$ ,  $\text{roof}(v_i)$  is a directed path in  $\mathcal{F}$  from  $x$  to  $y$  containing  $v_i$ , where the sub-path ending at  $v_i$  is  $\text{Lroof}(v_i)$ , and the sub-path starting  $v_i$  is  $\text{Rroof}(v_i)$ .

Let  $S$  be the set of vertices that comprise a largest antichain in  $\prec_{\mathcal{F}}$ , as illustrated in Fig. 2b with squares. Now consider the given geometric graph  $G$ . We may assume, by a suitable rotation, that no pair of vertices of  $G$  have the same  $y$ -coordinate. Let  $R$  be a largest ordered subset of  $S$  such that the  $y$ -coordinates of the vertices of  $R$  are either monotonically increasing or monotonically decreasing when considered in the order given by  $\sigma$ . By the Erdős–Szekeres Theorem [5],  $|R| \geq \sqrt{|S|}$ . Without loss of generality, assume that  $R$  is monotonically increasing. In what follows, we untangle  $G$  while keeping  $R$  fixed.

Let  $\mathcal{H}$  be the graph induced in  $\mathcal{E}$  by the following set of vertices:  $V(\mathcal{H}) := \{x, y\} \cup \{\text{roof}(w) : w \in R\}$ ; that is,  $\mathcal{H} = \mathcal{E}[V(\mathcal{H})]$ . Note that  $\mathcal{H}$  is not necessarily a subgraph of  $\mathcal{F}$ , as illustrated in Fig. 3.

We say that a simple polygonal chain  $C$  is *strictly x-monotone* if, for every vertical line  $\ell$ ,  $|C \cap \ell| \leq 1$ . For two distinct points  $p$  and  $q$  in the plane, let  $\overline{pq}$  denote the *open* line-segment with endpoints  $p$  and  $q$ . A simple polygon  $C$  is *star-shaped* (from  $p$ ) if there is a point  $p$  such that for every point  $q \in C$ ,  $\overline{pq} \cap C = \emptyset$ . The following lemma is the main ingredient in the proof of Lemma 4.

**Lemma 6** *The geometric planar graph  $G[V(\mathcal{H})]$  can be untangled so that each vertex of  $R$  is on the  $y$ -axis and it has the same  $y$ -coordinate in the untangling as in  $G[V(\mathcal{H})]$ . Moreover, all the internal faces of the untangling are star-shaped, and the path on its outer face from  $x$  to  $y$  not containing  $xy$  is strictly  $x$ -monotone.*



is a cycle and all internal faces are triangles. Thus that graph is a 2-connected graph, which provides the contradiction.

Applying Theorem 4 to embed each subgraph  $G^j$  yields an untangling of  $G$  in which the vertices of  $R$  are all on the  $y$ -axis and have their  $y$ -coordinates preserved. Applying Lemma 1 to this untangling completes the proof of the theorem.  $\square$

All that remains is to prove Lemma 6.


*Proof of Lemma 6* The proof is by induction on the number of vertices in  $R$ . We start by considering some useful properties of the roofs of two vertices in  $R$ .

Consider two vertices,  $u$  and  $v$  in  $R$  (or any two incomparable vertices in  $\langle \mathcal{F} \rangle$ , where  $u <_{\sigma} v$ ). Let  $x'$  be a vertex of  $\mathcal{F}$  such that  $x' \in \text{Lroof}(u)$  and  $x' \in \text{Lroof}(v)$  and such that the vertex following  $x'$  in  $\text{Lroof}(u)$  is not the same as the vertex following  $x'$  in  $\text{Lroof}(v)$ , as illustrated in Fig. 4. Similarly, let  $y'$  be a vertex of  $\mathcal{F}$  such that  $y' \in \text{Rroof}(u)$  and  $y' \in \text{Rroof}(v)$  and such that the vertex before  $y'$  in  $\text{Rroof}(u)$  is not the same as the vertex before  $y'$  in  $\text{Rroof}(v)$ . Such vertices,  $x'$  and  $y'$ , exist since  $u$  and  $v$  are incomparable in  $\mathcal{F}$ . Then  $\text{roof}(v)$  and  $\text{roof}(u)$  have the following properties. The paths between  $x$  and  $x'$  in  $\text{roof}(u)$  and in  $\text{roof}(v)$  coincide in  $\mathcal{F}$ , that is, the two paths are both equal to  $\text{Lroof}(x')$ . Similarly, the paths between  $y'$  and  $y$  in  $\text{roof}(u)$  and in  $\text{roof}(v)$  coincide in  $\mathcal{F}$ , that is, they are both equal to  $\text{Rroof}(y')$ . The path between  $x'$  and  $y'$  in  $\text{roof}(u)$  contains  $u$ , the path between  $x'$  and  $y'$  in  $\text{roof}(v)$  contains  $v$ , and the two paths have only  $x'$  and  $y'$  in common. Finally,  $u$  is inside the cycle determined by  $\text{roof}(v)$  and edge  $xy$  in  $\mathcal{F}$ . To summarize, for all  $u, v \in R$ , if  $u <_{\sigma} v$ , then each vertex of  $\text{roof}(u)$  is either on or inside the cycle determined by  $\text{roof}(v)$  and edge  $xy$  in  $\mathcal{F}$ .

We proceed by induction on the number of vertices in  $R$  but require a somewhat stronger inductive hypothesis than the statement of the lemma. Let  $C$  be a simple strictly  $x$ -monotone polygonal chain. We say that  $C$  is  $\varepsilon$ -ray-monotone from a point  $p = (x_p, y_p)$  if, for every point  $r = (x_p, y_p + t)$  with  $t \geq \varepsilon$  and every point  $q \in C$ ,  $(\overline{rq}) \cap C = \emptyset$ . Informally,  $C$  is  $\varepsilon$ -ray-monotone from  $p$  if every point sufficiently far above  $p$  sees all of  $C$ . Note that, under this definition, if  $C$  is  $\varepsilon$ -ray-monotone from  $p$ , then  $C$  is  $\varepsilon$ -ray-monotone from any point  $q = (x_p, y_p + t)$ ,  $t > 0$ , above  $p$ . Furthermore, there exists a value  $\delta = \delta(p, C, \varepsilon)$  such that  $C$  is  $\varepsilon$ -ray-monotone from any point  $p'$  whose distance from  $p$  is at most  $\delta$ . (This follows from the fact that the set of points  $p$  from which  $C$  is  $\varepsilon$ -ray-monotone is an open set.)

Let  $\varepsilon'$  be the minimum difference between the  $y$ -coordinates of any two vertices in  $R$ . We will construct a crossing-free geometric graph  $\overline{\mathcal{H}}$ , that is, an untangling

**Fig. 4** Roofs of two incomparable vertices  $u$  and  $v$  of  $\langle \mathcal{F} \rangle$



of  $G[V(\mathcal{H})]$ . In addition to the conditions of the lemma,  $\overline{\mathcal{H}}$  will have the following property: If  $|R| > 0$ , then the outer face of  $\overline{\mathcal{H}}$  is bounded by the edge  $xy$  and a path  $C$  from  $x$  to  $y$  such that  $C \cap R = \{v\}$  for some vertex  $v \in R$ , and  $C$  is  $\varepsilon$ -ray-monotone from  $v$  for some  $\varepsilon < \varepsilon'$ .

The base case occurs when  $|R| = 0$ . Then  $\mathcal{H}$  consists of the single edge  $xy$ , which can be untangled by placing  $x$  at  $(-1, t)$  and  $y$  at  $(1, t)$ , where  $t$  is smaller than any  $y$ -coordinate in  $G$ . Clearly this crossing-free geometric graph satisfies the conditions of the lemma and the inductive hypothesis. Next, suppose that  $|R| \geq 1$  and let  $v$  be the largest vertex of  $R$  in the total order  $\sigma$ . Let  $\mathcal{H}'$  be the subgraph of  $\mathcal{H}$  induced by the vertices in  $\{x, y\} \cup \{\text{roof}(u) : u \in R \setminus v\}$ . By induction, we can untangle  $G[V(\mathcal{H}')]$  to obtain a crossing-free geometric graph  $\overline{\mathcal{H}'}$  that satisfies the inductive hypothesis and the conditions of the lemma. It remains to place  $v$  and the vertices of  $\text{roof}(v)$  that are not yet placed. As described above, these vertices form a path  $P$  that goes from some vertex  $x'$  of  $\mathcal{H}'$  to  $v$  to some vertex  $y'$  of  $\mathcal{H}'$ . Denote by  $P'$  the path between  $x'$  and  $y'$  in  $\mathcal{H}'$  not containing  $xy$ . Furthermore, let  $E$  be the set of edges  $e$  in  $\mathcal{H}$  such that either both endpoints of  $e$  are interior vertices of  $P$ , or one endpoint of  $e$  is an interior vertex of  $P$  and the other endpoint of  $e$  is a vertex of  $P'$ .

The conditions of the lemma specify the location of  $v$ . In particular,  $v$  is on the  $y$ -axis, with its  $y$ -coordinate equal to its  $y$ -coordinate in  $G$ . The inductive hypothesis guarantees that the vertex  $v$  and any point sufficiently close to  $v$  can see<sup>5</sup> all vertices of the outer face of  $\overline{\mathcal{H}'}$ . Finally, we note that, if  $|R| > 1$ , then directly below  $v$ , on the  $y$ -axis, is a vertex  $u \in R$ . The fact that  $u$  is on the  $y$ -axis and that the outer face of  $\overline{\mathcal{H}'}$  is strictly  $x$ -monotone implies that the  $x$ -coordinate of  $x'$  is less than 0 and that the  $x$ -coordinate of  $y'$  is greater than 0. (For the special case where  $x' = x$  and/or  $y' = y$ , the above statement is still true.)

Next we place the interior vertices of  $P$  to obtain the crossing-free geometric graph  $\overline{\mathcal{H}}$ . To do this, draw a unit circle  $c$ , containing  $v$ , whose center is on the  $y$ -axis and below  $v$ . Place all interior vertices of  $P$  on  $c$  and sufficiently close to  $v$  so that:

- (1) the path on the outer face of  $\overline{\mathcal{H}}$  from  $x$  to  $y$  not containing  $xy$  is strictly  $x$ -monotone
- (2) in  $\overline{\mathcal{H}} \setminus E$ , all interior vertices of  $P$  see all other vertices of  $P$
- (3) in  $\overline{\mathcal{H}} \setminus E$ , all interior vertices of  $P$  see all vertices of  $P'$ , and
- (4) the path on the outer face of  $\overline{\mathcal{H}}$  from  $x$  to  $y$  not containing  $xy$  is  $\varepsilon$ -ray-monotone from  $v$  for some  $\varepsilon < \varepsilon'$ .

The first condition can be achieved since  $x'$  and  $y'$  are to the left and right, respectively, of the  $y$ -axis. The second condition can be achieved since we are placing the interior vertices of  $P$  on a convex curve (a circle) as close to  $v$  as necessary. The third condition can be achieved since the upper chain of  $\overline{\mathcal{H}'}$  is  $\varepsilon$ -ray-monotone from  $u$  and hence also from  $v$ . The fourth condition can be achieved by the definition of  $\varepsilon$ -ray-monotonicity and the first condition.

Consider the path in  $\overline{\mathcal{H}'}$  from  $x$  to  $y$  not containing  $xy$  along the outer face of  $\overline{\mathcal{H}'}$ . This path is comprised of the same vertices and edges as a directed path from  $x$  to  $y$

---

<sup>5</sup>Given a geometric graph, we say that a point  $p$  in the plane sees a point  $q$  if  $(\overline{pq})$  does not intersect the graph.

in  $\mathcal{F}$ . Thus, by Lemma 5, the outer face of  $\overline{\mathcal{H}}$  has no outer chords in  $\overline{\mathcal{H}}$ . Therefore, an edge of  $\overline{\mathcal{H}}$  that is not an edge of  $\overline{\mathcal{H}'}$  is either an edge of  $P$  or an edge in  $E$ . Clearly  $\overline{\mathcal{H}} \setminus E$  is crossing-free by the construction. The edges of  $E$  are not involved in any crossings in  $\overline{\mathcal{H}}$  by Conditions (2) or (3) above and the fact that  $\mathcal{H}$  is a planar embedding. Thus  $\overline{\mathcal{H}}$  is crossing-free. The vertices in  $R$  are all on the  $y$ -axis and all have the same  $y$ -coordinates in  $G$  as in  $\overline{\mathcal{H}}$ . Conditions (1) (and (4)) imply that the path between  $x$  and  $y$  on the outer face of  $\overline{\mathcal{H}}$  is strictly  $x$ -monotone. It remains to show that the internal faces of  $\overline{\mathcal{H}}$  are star-shaped. The only new faces in  $\overline{\mathcal{H}}$  not present in  $\overline{\mathcal{H}'}$  are the faces having interior vertices of  $P$  on their boundary. However, Conditions (2) and (3) above imply that each such face is star-shaped from some interior vertex of  $P$ . This completes the proof of the lemma.  $\square$

## 5 Trees: Upper Bound

In this section we prove the following theorem.

**Theorem 5** *For every positive number  $n$  such that  $\sqrt{n}$  is an integer, there exists a geometric forest (of stars)  $G$  on  $n$  vertices such that  $\text{fix}(G) = 3(\sqrt{n} - 1)$ . That is,  $G$  cannot be untangled while keeping more than  $3(\sqrt{n} - 1)$  vertices fixed, and  $G$  can be untangled while keeping exactly that many vertices fixed.*

*Proof* We first define  $G$ . A  $k$ -star is a rooted tree on  $k + 1$  vertices one of which is the root and the rest of the vertices are leaves adjacent to that root.  $G$  is a forest on  $n$  vertices comprised of trees  $T_i$ ,  $1 \leq i \leq \sqrt{n}$ , where each  $T_i$  is a  $(\sqrt{n} - 1)$ -star. All the vertices of  $G$  lie on the  $x$ -axis. For each  $i$ , the vertices of  $T_i$  have the  $x$ -coordinates  $i, i + \sqrt{n}, \dots, i + \sqrt{n}(\sqrt{n} - 1)$ , where the vertex with the maximum  $x$ -coordinate is the root of  $T_i$ . This completes the description of  $G$ .

*Upper Bound* We first prove that  $\text{fix}(G) \leq 3\sqrt{n} - 3$ ; that is, we prove that  $G$  cannot be untangled while keeping more than  $3\sqrt{n} - 3$  vertices fixed. Let  $H$  be an untangling of  $G$  with  $\text{fix}(G)$  vertices fixed. Let  $\ell$  denote the number of fixed leaves and  $r$  the number of fixed roots. Let  $r'$  denote the number of fixed roots that are adjacent to a fixed leaf. Given the ordering of the vertices of  $G$  on the  $x$ -axis, it is clear that  $r' \leq 1$ .

Partition the set of free roots into two sets. Let  $A$  be the set containing the free roots that are on or above the  $x$ -axis in  $H$ . Let  $B$  be the set containing the free roots that are strictly below the  $x$ -axis in  $H$ . Our reason for this nonsymmetric definition of  $A$  and  $B$  is to avoid double counting, and not because free roots on the  $x$ -axis have any special meaning. The total number of roots of  $G$  is  $|A| + |B| + r$ .

Suppose that the number of fixed leaves with a neighbor (i.e., a parent) in  $A$  is at most  $\sqrt{n} - 2 + |A|$ , and similarly for the number of fixed leaves with a neighbor in  $B$ . As noted above, at most one fixed leaf can be adjacent to a fixed root, and thus  $\ell \leq 2\sqrt{n} - 4 + |A| + |B| + r'$ . Since  $\text{fix}(G) = \ell + r$ , we get  $\text{fix}(G) \leq 2\sqrt{n} - 4 + |A| + |B| + r' + r$ . Having  $|A| + |B| + r = \sqrt{n}$  further implies that  $\text{fix}(G) \leq 3\sqrt{n} - 4 + r'$ . Since  $r' \leq 1$ , we get the desired upper bound.

Thus to complete the proof of the upper bound it remains to prove that the number of fixed leaves with a neighbor in  $A$  is at most  $\sqrt{n} - 2 + |A|$ . The proof below has

no special case for the free roots that are on the  $x$ -axis, so the proof for the number of fixed leaves with a neighbor in  $B$  is analogous.


Partition the leaves of  $G$  into a set of blocks  $\{P_j : 1 \leq j \leq \sqrt{n} - 1\}$  such that  $P_1$  contains the first  $\sqrt{n}$  leaves on the  $x$ -axis,  $P_2$  the next  $\sqrt{n}$  leaves, and so on. More formally,  $P_j$  contains all the leaves with  $x$ -coordinate in the range  $[1 + (j - 1)\sqrt{n}, j\sqrt{n}]$ . Note that each block contains exactly one leaf from each star of  $G$ . There are  $\sqrt{n} - 1$  blocks, each containing  $\sqrt{n}$  vertices.

Define an auxiliary graph  $Q$  with vertex set  $V(Q) = A \cup \{p_j : 1 \leq j \leq \sqrt{n} - 1\}$ , where  $vp_j \in E(Q)$  precisely if  $v$  is a vertex of  $A$  and  $v$  has a fixed neighbor in block  $P_j$ . Thus  $Q$  is a bipartite graph, where one bipartition is precisely the set  $A$ . Note that  $|V(Q)| = |A| + \sqrt{n} - 1$ . Since each vertex of  $A$  has exactly one neighbor in each block, the number of fixed leaves whose parents are in  $A$  is precisely  $|E(Q)|$ . We now show that  $Q$  has no cycles. That will complete the proof of the upper bound since in that case  $|E(Q)| \leq |V(Q)| - 1 = |A| + \sqrt{n} - 2$ .

Assume for the sake of contradiction that  $Q$  has a cycle. Let  $C$  be a shortest cycle in  $Q$ . Every second vertex of  $C$  is a vertex of  $A$ . The remaining vertices of  $C$  correspond to blocks of leaves. Let  $C_H$  be the subset of  $V(H)$  containing all the roots in  $V(C) \cap A$  and such that, for each of those roots,  $C_H$  also contains all its fixed leaves contained in blocks  $P_j$  for which  $p_j$  is in  $C$ .


Consider the geometric graph  $H[C_H]$ . The fact that  $C$  is a (shortest) cycle and that each vertex in  $A$  has exactly one leaf in each block, implies that  $H[C_H]$  is a geometric forest of 2-stars, where the vertices in  $V(C) \cap A$  have degree 2 in  $H[C_H]$ , and each block  $P_j$  such that  $p_j \in V(C) \setminus A$  has precisely two fixed leaves in  $H[C_H]$ .

Since  $H$  is crossing-free, so is  $H[C_H]$ . Furthermore, since all the roots in  $C_H$  are on or above the  $x$ -axis and all the leaves of  $C_H$  are on the  $x$ -axis,  $H[C_H]$  is fully contained in a closed half-plane determined by the  $x$ -axis. We now show that  $H[C_H]$  cannot be crossing-free, which will provide the desired contradiction.  $H[C_H]$  is a crossing-free geometric forest of 2-stars. We first expand  $H[C_H]$  into a crossing-free geometric cycle by adding some segments to it as follows. Consider blocks that contain a leaf of  $H[C_H]$ . Each such block  $P_j$  contains exactly two leaves of  $H[C_H]$ , denoted by  $j_1$  and  $j_2$  (see Fig. 5). We claim that  $\overline{j_1 j_2} \cap H[C_H] = \emptyset$ . There is no edge of  $H[C_H]$  that properly crosses  $\overline{j_1 j_2}$ , since  $H[C_H]$  is fully contained in a closed half-plane determined by the  $x$ -axis. Therefore,  $\overline{j_1 j_2} \cap H[C_H]$  can be nonempty only if there is an edge of  $H[C_H]$  fully contained in  $\overline{j_1 j_2}$ . This implies that there is a root of  $H[C_H]$  that is located on the  $x$ -axis between  $j_1$  and  $j_2$ . This however is impossible, since one of the two edges of  $H[C_H]$  incident to that root would contain  $j_1$  or  $j_2$  in its interior. This observation implies that  $H[C_H]$  can be extended into a




**Fig. 5** (Color online) Two 2-stars (one depicted in *green* and the other in *red*) with leaves in a common block

**Fig. 6** Illustration for the proof of the upper bound of Theorem 5



**Fig. 7** Untangled forest  $G$  with  $3\sqrt{n} - 3$  vertices fixed ( $k = \sqrt{n}$ )



*crossing-free* geometric cycle  $R$  by adding the appropriate line segments into each block that contains a leaf of  $H[C_H]$ .

Let  $v$  be, among all the roots in  $C_H$ , the one with the smallest index; that is, there is no other root  $w \in C_H$  where  $v \in T_i$  and  $w \in T_j$  with  $j < i$ . Vertex  $v$  has two neighbors (fixed leaves) in  $H[C_H]$ ,  $s_1 \in P_s$  and  $t_1 \in P_t$  (see Fig. 6). Vertex  $s_1$  has two neighbors in  $R$ . One is  $v$ , and the other is a vertex (fixed leaf)  $s_2 \in P_s$ . Similarly,  $t_1$  is adjacent in  $R$  to  $v$  and to a vertex (fixed leaf)  $t_2 \in P_t$ . Therefore,  $R$  contains two vertex disjoint paths:  $R_1$ , between  $s_1$  and  $t_1$ , and  $R_2$ , between  $s_2$  and  $t_2$ . Since  $v$  belongs to the smallest indexed tree, the ordering of their endpoints on the  $x$ -axis is  $s_1 < s_2 < t_1 < t_2$ . With such ordering of endpoints and since  $R$  is fully contained in the closed half-plane above the  $x$ -axis, it is impossible to draw  $R_1$  and  $R_2$  without crossings (since  $R_1$  separates the closed half-plane above the  $x$ -axis into two components, one containing  $s_2$  and one containing  $t_2$ ). This is the desired contradiction.

*Lower Bound* We now prove that  $\text{fix}(G) \geq 3\sqrt{n} - 3$ , that is, we prove that  $G$  can be untangled while keeping  $3\sqrt{n} - 3$  vertices fixed. Keep the followings vertices of  $G$  fixed:

- (1) all the leaves of  $T_1$  and  $T_2$
- (2) all the vertices in the block  $P_{\sqrt{n}-1}$ , and
- (3) the root of  $T_{\sqrt{n}}$ .

Move the root of  $T_1$  to the half-plane above the  $x$ -axis and move the root of  $T_2$  to the half-plane below the  $x$ -axis. For all  $3 \leq i \leq \sqrt{n} - 1$ , move all the free vertices of  $T_i$  to a very small disk centered at the fixed leaf of  $T_i$ . Move all the free leaves of  $T_{\sqrt{n}}$  to a small disk centered at the root of  $T_{\sqrt{n}}$ . Clearly, this can be done so that the resulting geometric forest  $H$  is crossing-free, as illustrated in Fig. 7. The number of fixed vertices of  $H$  is  $2(\sqrt{n} - 1) + (\sqrt{n} - 2) + 1 = 3\sqrt{n} - 3$ , as claimed.  $\square$

### 6 Conclusions

Polynomial bounds are now known for untangling all classes of planar graphs. Tight bounds (up to a constant) are known for untangling trees and outerplanar graphs. Gaps

remain open for untangling geometric cycles, where the best known lower and upper bounds are  $\Omega(n^{2/3})$  and  $\mathcal{O}((n \log n)^{2/3})$ , and for geometric planar graphs, where the best known lower and upper bounds are  $\Omega(n^{1/4})$  and  $\mathcal{O}(\sqrt{n})$ .

**Acknowledgements** This research was initiated at the Bellairs Workshop on Computational Geometry for Geometric Reconfigurations, 1 to 9 February 2007. The authors are grateful to Godfried Toussaint for organizing the workshop and to the other workshop participants for providing a stimulating working environment. Thanks to the two referees who provided many helpful suggestions.

## References

1. Abellanas, M., Hurtado, F., Ramos, P.: Tolerancia de arreglos de segmentos. In: Proc. VI Encuentros de Geometría Computacional, pp. 77–84 (1995)
2. Cıbulka, J.: Untangling polygons and graphs. In: Topological and Geometric Graph Theory. Electronic Notes in Discrete Mathematics, vol. 31, pp. 207–211 (2008)
3. de Fraysseix, H., Pach, J., Pollac, R.: How to draw a planar graph on a grid. *Combinatorica* **10**(1), 41–51 (1990)
4. Dilworth, R.P.: A decomposition theorem for partially ordered sets. *Ann. Math.* **51**(2), 161–166 (1950)
5. Erdős, P., Szekeres, G.: A combinatorial problem in geometry. *Compos. Math.* **2**, 464–470 (1935)
6. Fáry, I.: On straight line representation of planar graphs. *Acta Univ. Szeged. Sect. Sci. Math.* **11**, 229–233 (1948)
7. Goac, X., Kratochvil, J., Okamoto, Y., Shin, C.-S., Wolff, A.: Moving vertices to make drawings plane. In: Proc. 15th International Symp. on Graph Drawing (GD'07). Lecture Notes in Computer Science, vol. 4875, pp. 101–112. Springer, Berlin (2008). Also in <http://arxiv.org/abs/0706.1002>
8. Hong, S.-H., Nagamochi, H.: Convex drawings of graphs with non-convex boundary constraints. *Discrete Appl. Math.* **156**(12), 2368–2380 (2008)
9. Kang, M., Pikhurko, O., Ravsky, A., Schacht, M., Verbitsky, O.: Obfuscated drawings of planar graphs (2008). <http://arxiv.org/abs/0803.0858>
10. Kang, M., Schacht, M., Verbitsky, O.: How much work does it take to straighten a plane graph out? (2007). <http://arxiv.org/abs/0707.3373>
11. Pach, J., Tardos, G.: Untangling a polygon. *Discrete Comput. Geom.* **28**(4), 585–592 (2002)
12. Ramos, P.: Tolerancia de estructuras geométricas y combinatorias. Ph.D. thesis, Universidad Politécnica de Madrid, Madrid, Spain (1995)
13. Ravsky, A., Verbitsky, O.: On collinear sets in straight line drawings (2008). <http://arxiv.org/abs/0806.0253>
14. Spillner, A., Wolff, A.: Untangling a planar graph. In: Proc. 34th Internat. Conf. on Current Trends in Theory and Practice of Computer Science (SOFSEM'08). Lecture Notes in Computer Science, vol. 4910, pp. 473–484. Springer, Berlin (2008). Also in <http://arxiv.org/abs/0709.0170>
15. Verbitsky, O.: On the obfuscation complexity of planar graphs. *Theor. Comput. Sci.* **396**(1–3), 294–300 (2008)
16. Wagner, K.: Über eine Eigenschaft der ebene Komplexe. *Math. Ann.* **114**, 570–590 (1937)