

LAYERED SEPARATORS IN MINOR-CLOSED FAMILIES WITH APPLICATIONS

Vida Dujmović[†] Pat Morin[‡] David R. Wood[§]

Abstract. Graph separators are a ubiquitous tool in graph theory and computer science. However, in some applications, their usefulness is limited by the fact that the separator can be as large as $\Omega(\sqrt{n})$ in graphs with n vertices. This is the case for planar graphs, and more generally, for proper minor-closed families. We study a special type of graph separator, called a *layered separator*, which may have linear size in n , but has bounded size with respect to a different measure, called the *width*. We prove, for example, that planar graphs and graphs of bounded Euler genus admit layered separators of bounded width. More generally, we characterise the minor-closed classes that admit layered separators of bounded width as those that exclude a fixed apex graph as a minor.

We use layered separators to prove $\mathcal{O}(\log n)$ bounds for a number of problems where $\mathcal{O}(\sqrt{n})$ was a long standing previous best bound. This includes the *nonrepetitive chromatic number* and *queue-number* of graphs with bounded Euler genus. We extend these results to all proper minor-closed families, with a $\mathcal{O}(\log n)$ bound on the nonrepetitive chromatic number, and a $\log^{\mathcal{O}(1)} n$ bound on the queue-number. Only for planar graphs were $\log^{\mathcal{O}(1)} n$ bounds previously known. Our results imply that every graph from a proper minor-closed class has a *3-dimensional grid drawing* with $n \log^{\mathcal{O}(1)} n$ volume, whereas the previous best bound was $\mathcal{O}(n^{3/2})$.

June 7, 2013. Revised: April 15, 2015

[†] School of Computer Science and Electrical Engineering, University of Ottawa, Ottawa, Canada (vida.dujmovic@uottawa.ca). Research supported by NSERC and the Ontario Ministry of Research and Innovation.

[‡] School of Computer Science, Carleton University, Ottawa, Canada (morin@scs.carleton.ca). Research supported by NSERC

[§] School of Mathematical Sciences, Monash University, Melbourne, Australia (david.wood@monash.edu). Research supported by the Australian Research Council.

A short version of this paper and reference [22] was presented at the 54th Annual Symposium on Foundations of Computer Science (FOCS '13).

Contents

1	Introduction	3
1.1	Layered Separations	3
1.2	Queue-Number and 3-Dimensional Grid Drawings	5
1.3	Nonrepetitive Graph Colourings	6
2	Tree Decompositions and Separations	7
3	Graphs on Surfaces	9
4	Clique Sums	13
5	The Graph Minor Structure Theorem	14
6	Rich Decompositions and Shadow Complete Layerings	19
7	Track and Queue Layouts	21
8	3-Dimensional Graph Drawing	23
9	Nonrepetitive Colourings	24
10	Reflections	26
A	Separators in a Tree Decomposition	32
B	Layered Treewidth to Treewidth	33
C	Recursive Separators	33
D	Track Layout Construction	36

1 Introduction

Graph separators are a ubiquitous tool in graph theory and computer science since they are key to many divide-and-conquer and dynamic programming algorithms. Typically, the smaller the separator the better the results obtained. For instance, many problems that are $\mathcal{NP}$ -complete for general graphs have polynomial time solutions for classes of graphs that have bounded size separators—that is, graphs of bounded treewidth.

By the classical result of Lipton and Tarjan [49], every n -vertex planar graph has a separator of size $\mathcal{O}(\sqrt{n})$. More generally, the same is true for all proper minor-closed classes¹, as proved by Alon et al. [3]. While these results have found widespread use, separators of size $\Theta(\sqrt{n})$, or non-constant separators in general, are not small enough to be useful in some applications.

In this paper we study a type of graph separator, called layered separators, that may have $\Omega(n)$ vertices but have bounded size with respect to a different measure. In particular, layered separators intersect each layer of some predefined vertex layering in a bounded number of vertices. We prove that many classes of graphs admit such separators, and we show how they can be used to obtain logarithmic bounds for a variety of applications for which $\mathcal{O}(\sqrt{n})$ was the best known long-standing bound. These applications include nonrepetitive graph colourings, track layouts, queue layouts and 3-dimensional grid drawings of graphs. In addition, layered separators lend themselves to simple proofs.

In the remainder of the introduction, we define layered separators, and describe our results on the classes of graphs that admit them. Following that, we describe the implications that these results have on the above-mentioned applications.

1.1 Layered Separations

A *layering* of a graph G is a partition $(V_0, V_1, \dots, V_t)$ of $V(G)$ such that for every edge $vw \in E(G)$, if $v \in V_i$ and $w \in V_j$, then $|i - j| \leq 1$. Each set V_i is called a *layer*. For example, for a vertex r of a connected graph G , if V_i is the set of vertices at distance i from r , then $(V_0, V_1, \dots)$ is a layering of G , called the *bfs layering* of G starting from r . A *bfs tree* of G rooted at r is a spanning tree of G such that for every vertex v of G , the distance between v and r in G equals the distance between v and r in T . Thus, if $v \in V_i$ then the vr -path in T contains exactly one vertex from layer V_j for $j \in \{0, \dots, i\}$.

A *separation* of a graph G is a pair (G_1, G_2) of subgraphs of G such that $G = G_1 \cup G_2$. In particular, there is no edge between $V(G_1) - V(G_2)$ and $V(G_2) - V(G_1)$. The *order* of a separation (G_1, G_2) is $|V(G_1 \cap G_2)|$.

¹ A graph H is a *minor* of a graph G if a graph isomorphic to H can be obtained from a subgraph of G by contracting edges. A class $\mathcal{G}$ of graphs is *minor-closed* if $H \in \mathcal{G}$ for every minor H of G for every graph $G \in \mathcal{G}$. A minor-closed class is *proper* if it is not the class of all graphs.

A graph G admits *layered separations of width ℓ* with respect to a layering $(V_0, V_1, \dots, V_t)$ of G if for every set $S \subseteq V(G)$, there is a separation (G_1, G_2) of G such that:

- for $i \in \{0, 1, \dots, t\}$, layer V_i contains at most ℓ vertices in $V(G_1 \cap G_2)$, and
- both $V(G_1) - V(G_2)$ and $V(G_2) - V(G_1)$ contain at most $\frac{2}{3}|S|$ vertices in S .

Here the set $V(G_1 \cap G_2)$ is called a *layered separator of width ℓ* of $G[S]$. Note that these separators do not necessarily have small order, in particular $V(G_1 \cap G_2)$ can have $\Omega(n)$ vertices.

Layered separations are implicit in the seminal work of Lipton and Tarjan [49] on separators in planar graphs, and in many subsequent papers (such as [1, 39]). This definition was first made explicit by Dujmović et al. [23], who showed that a result of Lipton and Tarjan [49] implies that every planar graph admits layered separations of width 2. This result was used by Lipton and Tarjan as a subroutine in their $\mathcal{O}(\sqrt{n})$ separator result. We generalise this result for planar graphs to graphs embedded on arbitrary surfaces² In particular, we prove that graphs of Euler genus g admit layered separations of width $\mathcal{O}(g)$ (Theorem 10 in Section 3). A key to this proof is the notion of a layered tree decomposition, which is of independent interest, and is introduced in Section 2.

We further generalise this result by exploiting Robertson and Seymour’s graph minor structure theorem. Roughly speaking, a graph G is almost embeddable in a surface Σ if by deleting a bounded number of ‘apex’ vertices, the remaining graph can be embedded in Σ , except for a bounded number of ‘vortices’, where crossings are allowed in a well-structured way; see Section 5 where all these terms are defined. Robertson and Seymour proved that every graph from a proper minor-closed class can be obtained from clique-sums of graphs that are almost embeddable in a surface of bounded Euler genus. Here, apex vertices can be adjacent to any vertex in the graph. However, such freedom is not possible for graphs that admit layered separations of bounded width. For example, the planar $\sqrt{n} \times \sqrt{n}$ grid plus one dominant vertex (adjacent to every other vertex) does not admit layered separations of width $o(\sqrt{n})$ (see Section 5). We define the notion of strongly almost embeddable graphs, in which apex vertices are only allowed to be adjacent to vortices and other apex vertices. With this restriction, we prove that graphs obtained from clique-sums of strongly almost embeddable graphs admits layered separations of bounded width (Theorem 19 in Section 5). A recent structure theorem of Dvořák and Thomas [34] says that H -minor-free graphs have this structure, for each apex graph H . (A graph H is *apex* if $H - v$ is planar for some vertex v .) We conclude that a minor-closed class $\mathcal{G}$ admits layered separations of bounded width if and only if $\mathcal{G}$ excludes some fixed apex graph. Then, in all the applications that we consider, we deal with (unrestricted) apex vertices separately, leading to $\mathcal{O}(\log n)$ or $\log^{\mathcal{O}(1)} n$ bounds for every proper minor-closed family. These extensions depend on

² The *Euler genus* of a surface Σ is $2 - \chi$, where χ is the Euler characteristic of Σ . Thus the orientable surface with h handles has Euler genus $2h$, and the non-orientable surface with c cross-caps has Euler genus c . The *Euler genus* of a graph G is the minimum Euler genus of a surface in which G embeds. See [51] for background on graphs embedded in surfaces.

two tools of independent interest (rich tree decompositions and shadow complete layerings) that are presented in Section 6.

1.2 Queue-Number and 3-Dimensional Grid Drawings

Let G be a graph. In a linear ordering $\preceq$ of $V(G)$, two edges vw and xy are *nested* if $v \prec x \prec y \prec w$. A k -*queue layout* of a graph G consists of a linear ordering $\preceq$ of $V(G)$ and a partition $E_1, \dots, E_k$ of $E(G)$, such that no two edges in each set E_i are nested with respect to $\preceq$. The *queue-number* of a graph G is the minimum integer k such that G has a k -queue layout, and is denoted by $\text{qn}(G)$. Queue layouts were introduced by Heath and Rosenberg [45, 46] and have since been widely studied. They have applications in parallel process scheduling, fault-tolerant processing, matrix computations, and sorting networks; see [29, 55] for surveys.

A number of classes of graphs are known to have bounded queue-number. For example, every tree has a 1-queue layout [46], every outerplanar graph has a 2-queue layout [45], every series-parallel graph has a 3-queue layout [57], every graph with bandwidth b has a $\lceil \frac{b}{2} \rceil$ -queue layout [46], every graph with pathwidth p has a p -queue layout [26], and more generally every graph with bounded treewidth has bounded queue-number [26]. All these classes have bounded treewidth. Only a few highly structured graphs of unbounded treewidth, such as grids and cartesian products [69], are known to have bounded queue-number. In particular, it is open whether planar graphs have bounded queue-number, as conjectured by Heath *et al.* [45, 46].

The dual concept of a queue layout is a *stack layout*, introduced by Ollmann [53] and commonly called a *book embedding*. It is defined similarly, except that no two edges in the same set are allowed to cross with respect to the vertex ordering. *Stack-number* (also known as *book thickness* or *page-number*) is bounded for planar graphs [73], for graphs of bounded Euler genus [50], and for every proper minor-closed graph family [7]. A recent construction of bounded degree monotone expanders by Bourgain [9, 10] has bounded stack-number and bounded queue-number; see [25, 28, 33].

Until recently, the best known upper bound for the queue-number of planar graphs was $\mathcal{O}(\sqrt{n})$. This upper bound follows easily from the fact that planar graphs have pathwidth at most $\mathcal{O}(\sqrt{n})$. In a breakthrough result, this bound was reduced to $\mathcal{O}(\log^2 n)$ by Di Battista, Frati, and Pách [18], which was further improved by Dujmović [22] to $\mathcal{O}(\log n)$ using a simple proof based on layered separators. In particular, Dujmović [22] proved that every n -vertex graph that admits layered separations of width ℓ has $\mathcal{O}(\ell \log n)$ queue-number. Since every planar graph admits layered separations of width 2, planar graphs have $\mathcal{O}(\log n)$ queue-number [22]. Moreover, we immediately obtain logarithmic bounds on the queue-number for the graph families described in Section 1.1. In particular, we prove that graphs with Euler genus g have $\mathcal{O}(g \log n)$ queue-number (Theorem 27), and graphs that exclude a fixed apex minor have $\mathcal{O}(\log n)$ queue-number (Theorem 28). Furthermore, we extend this result to all proper minor-closed families with an upper bound of $\log^{\mathcal{O}(1)} n$ (Theorem 30). The previously best known bound for all these families,

except for planar graphs, was $\mathcal{O}(\sqrt{n})$.

One motivation for studying queue layouts is their connection with 3-dimensional graph drawing. A *3-dimensional grid drawing* of a graph G represents the vertices of G by distinct grid points in $\mathbb{Z}^3$ and represents each edge of G by the open segment between its endpoints, such that no two edges intersect. The *volume* of a 3-dimensional grid drawing is the number of grid points in the smallest axis-aligned grid-box that encloses the drawing. For example, Cohen et al. [13] proved that the complete graph K_n has a 3-dimensional grid drawing with volume $\mathcal{O}(n^3)$ and this bound is optimal. Pach et al. [54] proved that every graph with bounded chromatic number has a 3-dimensional grid drawing with volume $\mathcal{O}(n^2)$, and this bound is optimal for $K_{n/2, n/2}$. More generally, Bose et al. [8] proved that every 3-dimensional grid drawing of an n -vertex m -edge graph has volume at least $\frac{1}{8}(n + m)$. Dujmović and Wood [30] proved that every graph with bounded maximum degree has a 3-dimensional grid drawing with volume $\mathcal{O}(n^{3/2})$, and the same bound holds for graphs from a proper minor-closed class. In fact, every graph with bounded degeneracy has a 3-dimensional grid drawing with $\mathcal{O}(n^{3/2})$ volume [32]. Dujmović et al. [26] proved that every graph with bounded treewidth has a 3-dimensional grid drawing with volume $\mathcal{O}(n)$. Whether planar graphs have 3-dimensional grid drawings with $\mathcal{O}(n)$ volume is a major open problem, due to Felsner et al. [38]. The best known bound on the volume of 3-dimensional grid drawings of planar graphs is $\mathcal{O}(n \log n)$ by Dujmović [22]. We prove a $\mathcal{O}(n \log n)$ volume bound for graphs of bounded Euler genus (Theorem 32), and more generally, for apex-minor-free graphs (Theorem 33). Most generally, we prove an $n \log^{\mathcal{O}(1)} n$ volume bound for every proper minor-closed family (Theorem 34).

All our results about queue layouts are proved in Section 7, and all our results about 3-dimensional grid drawings are proved in Section 8.

1.3 Nonrepetitive Graph Colourings

A vertex colouring of a graph is *nonrepetitive* if there is no path for which the first half of the path is assigned the same sequence of colours as the second half. More precisely, a k -colouring of a graph G is a function ψ that assigns one of k colours to each vertex of G . A path $(v_1, v_2, \dots, v_{2t})$ of even order in G is *repetitively* coloured by ψ if $\psi(v_i) = \psi(v_{t+i})$ for $i \in \{1, \dots, t\}$. A colouring ψ of G is *nonrepetitive* if no path of G is repetitively coloured by ψ . Observe that a nonrepetitive colouring is *proper*, in the sense that adjacent vertices are coloured differently. The *nonrepetitive chromatic number* $\pi(G)$ is the minimum integer k such that G admits a nonrepetitive k -colouring.

The seminal result in this area is by Thue [65], who in 1906 proved that every path is non-repetitively 3-colourable. Nonrepetitive colourings have recently been widely studied; see the surveys [12, 41, 42]. A number of graph classes are known to have bounded nonrepetitive chromatic number. In particular, trees are nonrepetitively 4-colourable [11, 48], outerplanar graphs are nonrepetitively 12-colourable [5, 48], and more generally, every graph with treewidth k is nonrepetitively 4^k -colourable [48]. Graphs with maximum degree Δ are nonrepetitively $\mathcal{O}(\Delta^2)$ -

colourable [2, 24, 41, 44].

Perhaps the most important open problem in the field of nonrepetitive colourings is whether planar graphs have bounded nonrepetitive chromatic number [2]. The best known lower bound is 11, due to Ochem [23]. Dujmović et al. [23] showed that layered separations can be used to construct nonrepetitive colourings. In particular, every n -vertex graph that admits layered separations of width ℓ is nonrepetitively $\mathcal{O}(\ell \log n)$ -colourable [23]. Applying the result for planar graphs mentioned above, Dujmović et al. [23] concluded that every n -vertex planar graph is nonrepetitively $\mathcal{O}(\log n)$ -colourable. We generalise this result to conclude that every graph with Euler genus g is nonrepetitively $\mathcal{O}(g + \log n)$ -colourable (Theorem 38). The previous best bound for graphs of bounded genus was $\mathcal{O}(\sqrt{n})$, which is obtained by an easy application of the standard $\mathcal{O}(\sqrt{n})$ separator result for graphs of bounded genus. We further generalise this result to conclude a $\mathcal{O}(\log n)$ bound for arbitrary proper minor-closed classes (Theorem 40).

All our results about nonrepetitive graph colouring are proved in Section 9.

2 Tree Decompositions and Separations

Graphs decompositions, especially tree decompositions, are a key to our results. For graphs G and H , an H -decomposition of G is a collection $(B_x \subseteq V(G) : x \in V(H))$ of sets of vertices in G (called *bags*) indexed by the vertices of H , such that:

- (1) for every edge vw of G , some bag B_x contains both v and w , and
- (2) for every vertex v of G , the set $\{x \in V(H) : v \in B_x\}$ induces a non-empty connected subgraph of H .

The *width* of a decomposition is the size of the largest bag minus 1. If H is a tree, then an H -decomposition is called a *tree decomposition*. The *treewidth* of a graph G is the minimum width of any tree decomposition of G . Tree decompositions were first introduced by Halin [43] and independently by Robertson and Seymour [60]. H -decompositions, for general graphs H , were introduced by Diestel and Kühn [20]; also see [72].

Separators and treewidth are closely connected, as shown by the following two results (see Appendix A).

Lemma 1 (Robertson and Seymour [60], (2.5) & (2.6)). *If S is a set of vertices in a graph G , then for every tree decomposition of G there is a bag B such that each connected component of $G - B$ contains at most $\frac{1}{2}|S|$ vertices in S , which implies that G has a separation (G_1, G_2) with $V(G_1 \cap G_2) = B$ and both $G_1 - V(G_2)$ and $G_2 - V(G_1)$ contain at most $\frac{2}{3}|S|$ vertices in S .*

Lemma 2 (Reed [56], Fact 2.7). *Assume that for every set S of vertices in a graph G , there is a separation (G_1, G_2) of G such that $|V(G_1 \cap G_2)| \leq k$ and both $V(G_1) - V(G_2)$ and $V(G_2) - V(G_1)$ contain at most $\frac{2}{3}|S|$ vertices in S . Then G has treewidth less than $4k$.*

We now define the *layered width* of a decomposition, which is the key original definition of this paper. The *layered width* of an H -decomposition $(B_x : x \in V(H))$ of a graph G is the minimum integer ℓ such that, for some layering $(V_0, V_1, \dots, V_t)$ of G , each bag B_x contains at most ℓ vertices in each layer V_i . The *layered treewidth* of a graph G is the minimum layered width of a tree decomposition of G . Note that if we only consider layerings in which every vertex is in a single layer, then layered treewidth equals treewidth plus 1.

The following result, which is implied by Lemma 1, shows that bounded layered treewidth leads to layered separations of bounded width; see Theorem 21 for a converse result.

Lemma 3. *Every graph with layered treewidth ℓ admits layered separations of width at most ℓ .*

The *diameter* of a connected graph G is the maximum distance of two vertices in G . Layered tree decompositions lead to tree decompositions of bounded width for graphs of bounded diameter.

Lemma 4. *If a connected graph G has diameter d , treewidth k and layered treewidth ℓ , then $k < \ell(d + 1)$.*

Proof. Every layering of G has at most $d + 1$ layers. Thus each bag in a tree decomposition of layered width ℓ contains at most $\ell(d + 1)$ vertices. The claim follows. $\square$

Similarly, a graph of bounded diameter that admits layered separations of bounded width has bounded treewidth.

Lemma 5. *If a connected graph G has diameter d , treewidth k and admits layered separations of width ℓ , then $k < 4\ell(d + 1)$.*

Proof. Since G admits layered separations of width ℓ , there is a layering of G such that for every set $S \subseteq V(G)$, there is a separation (G_1, G_2) of G such that each layer contains at most ℓ vertices in $V(G_1 \cap G_2)$, and both $V(G_1) - V(G_2)$ and $V(G_2) - V(G_1)$ contain at most $\frac{2}{3}|S|$ vertices in S . Since G has diameter d , the number of layers is at most $d + 1$. Thus $|V(G_1 \cap G_2)| \leq (d + 1)\ell$. The claim follows from Lemma 2. $\square$

Lemmas 4 and 5 can essentially be rewritten in the language of ‘local treewidth’, which was first introduced by Eppstein [36] under the guise of the ‘treewidth–diameter’ property. A graph class $\mathcal{G}$ has *bounded local treewidth* if there is a function f such that for every graph G in $\mathcal{G}$, for every vertex v of G and for every integer $r \geq 0$, the subgraph of G induced by the vertices at distance at most r from v has treewidth at most $f(r)$; see [14, 16, 36, 40]. If $f(r)$ is a linear function, then $\mathcal{G}$ has *linear local treewidth*.

Lemma 6. *If every graph in some class $\mathcal{G}$ has layered treewidth at most ℓ , then $\mathcal{G}$ has linear local treewidth with $f(r) = \ell(2r + 1) - 1$.*

Proof. Given a vertex v in a graph $G \in \mathcal{G}$, and given an integer $r \geq 0$, let G' be the subgraph of G induced by the set of vertices at distance at most r from v . By assumption, G has a tree decomposition of layered width ℓ with respect to some layering $(V_0, V_1, \dots, V_t)$. If $v \in V_i$ then $V(G') \subseteq V_{i-r} \cup \dots \cup V_{i+r}$. Thus G' contains at most $(2r+1)\ell$ vertices in each bag. Hence G has treewidth at most $(2r+1)\ell - 1$, and $\mathcal{G}$ has linear local treewidth. $\square$

Lemma 7. *If every graph in some class $\mathcal{G}$ admits layered separations of width at most ℓ , then $\mathcal{G}$ has linear local treewidth with $f(r) < 4\ell(2r+1)$.*

Proof. Given a vertex v in a graph $G \in \mathcal{G}$, and given an integer $r \geq 0$, let G' be the subgraph of G induced by the set of vertices at distance at most r from v . By assumption, there is a layering $(V_0, V_1, \dots, V_t)$ of G such that for every set $S \subseteq V(G)$, there is a separation (G_1, G_2) of G such that each layer contains at most ℓ vertices in $V(G_1 \cap G_2)$, and both $V(G_1) - V(G_2)$ and $V(G_2) - V(G_1)$ contain at most $\frac{2}{3}|S|$ vertices in S . If $v \in V_i$ then $V(G') \subseteq V_{i-r} \cup \dots \cup V_{i+r}$. Thus $|V(G_1 \cap G_2 \cap G')| \leq (2r+1)\ell$. By Lemma 2, G' has treewidth less than $4(2r+1)\ell$. The claim follows. $\square$

We conclude this section with a few observations about layered treewidth.

First we bound the number of edges in a graph G with layered treewidth k . Let S be a leaf bag in a tree decomposition of G with layered width k . Let T be the neighbouring bag. If $S \subseteq T$ then delete S and repeat. Otherwise there is a vertex v in $S \setminus T$. Say v is in layer V_i . Then every neighbour of v is in $S \cap (V_{i-1} \cup V_i \cup V_{i+1}) \setminus \{v\}$, which has size at most $3k-1$. Thus G has minimum degree at most $3k-1$, which implies that G has at most $(3k-1)n$ edges. The following example shows that this bound is roughly tight. For integers $p \gg k \geq 2$, let G be the graph with vertex set $\{(x, y) : x, y \in \{1, \dots, p\}\}$, where distinct vertices (x, y) and (x', y') are adjacent if $|y - y'| \leq 1$ and $|x - x'| \leq k-1$. For $y \in \{1, \dots, p\}$, let $V_y := \{(x, y) : x \in \{1, \dots, p\}\}$. Then $(V_1, V_2, \dots, V_p)$ is a layering of G . For $x \in \{1, \dots, p-k+1\}$, let $B_x := \{(x', y) : x' \in \{x, \dots, x+k-1\}, y \in \{1, \dots, p\}\}$. Then $B_1, B_2, \dots, B_{p-k+1}$ is a tree decomposition of G with layered width k . Apart from vertices near the boundary, every vertex of G has degree $6k-4$. Thus $|E(G)| = (3k-2)n - \mathcal{O}(k\sqrt{n})$.

Finally, note that layered treewidth is not a minor-closed parameter. For example, if G is the 3-dimensional $n \times n \times 2$ grid graph, then it is easily seen that G has layered treewidth at most 3, but G contains a K_n minor [71], and K_n has layered treewidth $\lceil \frac{n}{2} \rceil$.

3 Graphs on Surfaces

This section constructs layered tree decompositions of graphs with bounded Euler genus. The following definitions and simple lemma will be useful. A *triangulation* of a surface is a loopless

multigraph embedded in the surface, such that each face is bounded by three edges. We emphasise that parallel edges not bounding a single face are allowed. For a subgraph G' of G , let $F(G')$ be the set of faces of G incident with at least one vertex of G' . Let G^* be the dual of G . That is, $V(G^*) = F(G)$ where $fg \in E(G^*)$ whenever some edge of G is incident with both f and g (for all distinct faces $f, g \in F(G)$). Thus the edges of G are in 1–1 correspondence with the edges of G^* . Let T be a subtree of G . An edge $vw \in E(G)$ is a *chord* of T if $v, w \in V(T)$ and $vw \notin E(T)$. An edge $vw \in E(G)$ is a *half-chord* of T if $|\{v, w\} \cap V(T)| = 1$. An edge of G^* dual to a chord of G is called a *dual-chord*. An edge of G^* dual to a half-chord of G is called a *dual-half-chord*.

Lemma 8. *Let T be a non-empty subtree of a triangulation G of a surface. Let H be the subgraph of G^* with vertex set $F(T)$ and edge set the dual-chords and dual-half-chords of T . Then H is connected. Moreover, $H - e$ is connected for each dual-half-chord e of T .*

Proof. Consider the following walk W in T . Choose an arbitrary edge $\alpha\beta$ in T , and initialise $W := (\alpha, \beta)$. Apply the following rule to choose the next vertex in W . Suppose that $W = (\alpha, \beta, \dots, x, y)$. Let yz be the edge of T incident with y clockwise from yx in the cyclic ordering of edges incident to y defined by the embedding of T . (It is possible that $x = z$.) Then append z to W . Stop when the edge $\alpha\beta$ is traversed in this order for the second time. Clearly each edge of T is traversed by W exactly two times (once in each direction). Let W' be the walk in H obtained from W as follows. Consider three consecutive vertices x, y, z in W . Let $f_1, f_2, \dots, f_k$ be the sequence of faces from yx to yz in the clockwise order of faces incident with y . Construct W' from W by replacing y by $f_1, f_2, \dots, f_k$ (and doing this simultaneously at each vertex y in W). Each such face f_i is incident with y , and is thus a vertex of H . Moreover, for $i \in \{1, \dots, k-1\}$, the edge $f_i f_{i+1}$ of G^* is dual to a chord or half-chord of T , and thus $f_i f_{i+1}$ is an edge of H . Hence W' is a walk in H . Every face of G incident with at least one vertex in T appears in W' . Thus W' is a spanning walk in H . Therefore H is connected, as claimed.

Let H' be the subgraph of H formed by the dual-half-chords of T . We now show that H' is 2-regular. Consider a dual-half-chord fg of T . Let vw be the corresponding half-chord of G , where $v \in V(T)$ and $w \notin V(T)$. Say u is the third vertex incident to f . If $u \in V(T)$ then uv is not a half-chord of T and uw is a half-chord of T , implying that the only other neighbour of f in H' (in addition to g) is the other face incident to uw (in addition to f). On the other hand, if $u \notin V(T)$ then uv is a half-chord of T and uw is not a half-chord of T , implying that the only other neighbour of f in H' (in addition to g) is the other face incident to uv (in addition to f). Hence f has degree 2 in H' , and H' is 2-regular. Therefore, if e is a dual-half-chord of T , then e is in a cycle, and $H - e$ is connected. $\square$

The following theorem is the main result of this section.

Theorem 9. *Every graph G with Euler genus g has layered treewidth at most $2g + 3$.*




Figure 1: Construction of H in Lemma 8.

Proof. We may assume that G is a triangulation of a surface with Euler genus g . Let r be a vertex of G . Let $F(G)$ be the set of faces of G . Say G has n vertices. By Euler's formula, $|F(G)| = 2n + 2g - 4$ and $|E(G)| = 3n + 3g - 6$. Let $(V_0, V_1, \dots, V_t)$ be the bfs layering of G starting from r . Let T be a bfs tree of G rooted at r . For each vertex v of G , let P_v be the vertex set of the vr -path in T . Thus if $v \in V_i$, then P_v contains exactly one vertex in V_j for $j \in \{0, \dots, i\}$.

Let D be the subgraph of G^* with vertex set $F(G)$, where two vertices are adjacent if the corresponding faces share an edge not in T . Thus $|V(D)| = |F(G)| = 2n + 2g - 4$ and $|E(D)| = |E(G)| - |E(T)| = (3n + 3g - 6) - (n - 1) = 2n + 3g - 5$. Since $V(T) = V(G)$, each edge of G is either an edge of T or is a chord of T . Thus D is the graph H defined in Lemma 8. By Lemma 8, D is connected.

Let T^* be a spanning tree of D . Thus $|E(T^*)| = |V(D)| - 1 = 2n + 2g - 5$. Let $X := E(D) - E(T^*)$. Thus $|X| = (2n + 3g - 5) - (2n + 2g - 5) = g$. For each face $f = xyz$ of G , let

$$C_f := \cup\{P_a \cup P_b : ab \in X\} \cup P_x \cup P_y \cup P_z .$$

Since $|X| = g$ and each P_v contains at most one vertex in each layer, C_f contains at most $2g + 3$ vertices in each layer.

We claim that $(C_f : f \in F(G))$ is a T^* -decomposition of G . For each edge vw of G , if f is a face incident to vw then v and w are in C_f . This proves condition (1) in the definition of T^* -decomposition.

We now prove condition (2). It suffices to show that for each vertex v of G , if F' is the set of faces f of G such that v is in C_f , then the induced subgraph $T^*[F']$ is connected and non-empty. Each face incident to v is in F' , thus F' is non-empty. Let T' be the subtree of T rooted at v .

If some edge xy in X is a half-chord or chord of T' , then v is in $P_x \cup P_y$, implying that v is in every bag, and $T^*[F'] = T^*$ is connected. Now assume that no half-chord or chord of T' is in X . Thus a face f of G is in F' if and only if f is incident with a vertex in T' ; that is, $F' = F(T')$. If $v = r$, then $T' = T$ and $F' = F(G)$, implying $T^*[F'] = T^*$, which is connected. Now assume that $v \neq r$. Let p be the parent of v in T . Let H be the graph defined in Lemma 8 with respect to T' . So H has vertex set F' and edge set the dual-chords and dual-half-chords of T' . Each chord or half-chord of T' is an edge of $G - (E(T) \cup X)$, except for pv , which is a half-chord of T' (since $p \notin V(T')$). Let e be the edge of H dual to pv . By Lemma 8, $T^*[F'] = H - e$ is connected, as desired.

Therefore $(C_f : f \in F(G))$ is a T^* -decomposition of G with layered width at most $2g + 3$. $\square$

Several notes on Theorem 9 are in order.

- A spanning tree in an embedded graph with an ‘interdigitating’ spanning tree in the dual was introduced for planar graphs by von Staudt [67] in 1847, and is sometimes called a *tree-cotree decomposition* [37]. This idea was generalised for orientable surfaces by Biggs [6] and for non-orientable surfaces by Richter and Shank [58]; also see [64].
- Lemma 3 and Theorem 9 imply the following result for layered separators.

Theorem 10. *Every graph with Euler genus g admits layered separations of width $2g + 3$.*

Sergey Norin [personal communication, 2014] proved that every n -vertex graph G with layered treewidth k has treewidth at most $2\sqrt{kn}$ (see Appendix B). Thus Theorem 9 implies:

Theorem 11. *Every n -vertex graph with Euler genus g has treewidth at most $2\sqrt{(2g + 3)n}$.*

Lemma 1 then implies that n -vertex graphs of Euler genus g have separators of order $\mathcal{O}(\sqrt{gn})$, as proved in [1, 21, 37, 39]. Gilbert et al. [39] gave examples of such graphs with no $o(\sqrt{gn})$ separator, and thus with treewidth $\Omega(\sqrt{gn})$ (by Lemma 1). Hence each of the upper bounds in Theorem 9–11 are within a constant factor of optimal.

Note that the proof of Theorem 9 uses ideas from many previous proofs about separators in embedded graphs [1, 37, 39]. For example, Aleksandrov and Djidjev [1] call the graph D in the proof of Theorem 9 a *separation graph*.

- If we apply Theorem 9 to a graph with radius d , where r is a central vertex, then each bag consists of $2g + 3$ paths ending at r , each of length at most d . Thus each bag contains at most $(2g + 3)d + 1$ vertices. We obtain the following result, first proved in the planar case by Robertson and Seymour [59] and implicitly by Baker [4], and in general by Eppstein [36] with a $\mathcal{O}(gd)$ bound. Eppstein’s proof also uses the tree-cotree decomposition; see [35, 37] for related work.

Theorem 12. *Every graph with Euler genus g and radius d has treewidth at most $(2g+3)d$. In particular, every planar graph with radius d has treewidth at most $3d$.*

- The proof of Theorem 9 gives the following stronger result that will be useful later, where $Q = \cup\{P_a \cup P_b : ab \in X\}$.

Theorem 13. *Let r be a vertex in a graph G with Euler genus g . Then there is a tree decomposition $\mathcal{T}$ of G with layered width at most $2g+3$ with respect to some layering in which the first layer is $\{r\}$. Moreover, there is a set $Q \subseteq V(G)$ with at most $2g$ vertices in each layer, such that $\mathcal{T}$ restricted to $G - Q$ has layered width at most 3 with respect to the same layering.*

4 Clique Sums

We now extend the above results to more general graph classes via the clique-sum operation. For compatibility with this operation, we introduce the following concept that is slightly stronger than having bounded layered treewidth. A *clique* is a set of pairwise adjacent vertices in a graph. Say a graph G is ℓ -good if for every clique K of size at most ℓ in G there is a tree decomposition of G of layered width at most ℓ with respect to some layering of G in which K is the first layer.

Theorem 14. *Every graph G with Euler genus g is $(2g+3)$ -good.*

Proof. Given a clique K of size at most $2g+3$ in G , let G' be the graph obtained from G by contracting K into a single vertex r . Then G' has Euler genus at most g . Theorem 13 gives a tree decomposition of G' of layered width at most $2g+3$ with respect to some layering of G' in which $\{r\}$ is the first layer. Replace the first layer by K , and replace each instance of r in the tree decomposition of G' by K . We obtain a tree decomposition of G of layered width at most $2g+3$ with respect to some layering of G in which K is the first layer (since $|K| \leq 2g+3$). Thus G is $(2g+3)$ -good. $\square$

Let $C_1 = \{v_1, \dots, v_k\}$ be a k -clique in a graph G_1 . Let $C_2 = \{w_1, \dots, w_k\}$ be a k -clique in a graph G_2 . Let G be the graph obtained from the disjoint union of G_1 and G_2 by identifying v_i and w_i for $i \in \{1, \dots, k\}$, and possibly deleting some edges in $C_1 (= C_2)$. Then G is a k -clique sum of G_1 and G_2 . If $k \leq \ell$ then G is a $(\leq \ell)$ -clique sum of G_1 and G_2 .

Lemma 15. *For $\ell \geq k$, if G is a $(\leq k)$ -clique-sum of ℓ -good graphs G_1 and G_2 , then G is ℓ -good.*

Proof. Let K be the given clique of size at most ℓ in G . Without loss of generality, K is in G_1 . Since G_1 is ℓ -good, there is a tree decomposition T_1 of G_1 of layered width at most ℓ with respect to some layering of G_1 in which K is the first layer. Let $X := V(G_1 \cap G_2)$. Thus X is a clique in G_1 and in G_2 . Hence X is contained in at most two consecutive layers of the above layering of G_1 . Let X' be the subset of X in the first of these two layers. Note that if $K \cap X \neq \emptyset$

then $X' = K \cap X$. Since $|X'| \leq k \leq \ell$ and since G_2 is ℓ -good, there is a tree decomposition T_2 of G_2 with layered width at most ℓ with respect to some layering of G_2 in which X' is the first layer. Thus the second layer of G_2 contains $X \setminus X'$. Now, the layerings of G_1 and G_2 can be overlaid, with the layer containing X' in common, and the layer containing $X \setminus X'$ in common. By the definition of X' , it is still the case that the first layer is K . Let T be the tree decomposition of G obtained from the disjoint union of T_1 and T_2 by adding an edge between a bag in T_1 containing X and a bag in T_2 containing X . (Each clique is contained in some bag of a tree decomposition.) For each bag B of T the intersection of B with a single layer consists of the same set of vertices as the intersection of B and the corresponding layer in the layering of G_1 or G_2 . Hence T has layered width at most ℓ . $\square$

We now describe some graph classes for which Lemma 15 is immediately applicable. Wagner [68] proved that every K_5 -minor-free graph can be constructed from (≤ 3) -clique sums of planar graphs and V_8 , where V_8 is the graph obtained from an 8-cycle by adding an edge between opposite vertices. A bfs layering shows that V_8 is 3-good. By Theorem 14, every planar graph is 3-good. Thus, by Lemma 15, every K_5 -minor-free graph is 3-good. By Lemma 3, such a graph has layered treewidth at most 3, and admits layered separations of width 3. Wagner [68] also proved that every $K_{3,3}$ -minor-free graph can be constructed from (≤ 2) -clique sums of planar graphs and K_5 . Since K_5 is 4-good and every planar graph is 3-good, every $K_{3,3}$ -minor-free graph is 4-good, has layered treewidth at most 4, and admits layered separations of width 4. For a number of particular graphs H , Truemper [66] characterised the H -minor-free graphs in terms of (≤ 3) -clique sums of planar graphs and various small graphs. The above methods apply here also; we omit these details. More generally, a graph H is *single-crossing* if it has a drawing in the plane with at most one crossing. For example, K_5 and $K_{3,3}$ are single-crossing. Robertson and Seymour [61] proved that for every single-crossing graph H , every H -minor-free graph can be constructed from (≤ 3) -clique sums of planar graphs and graphs of treewidth at most ℓ , for some constant $\ell = \ell(H) \geq 3$. It follows from the above results that every H -minor-free graph is ℓ -good, has layered treewidth at most ℓ , and admits layered separations of width ℓ .

5 The Graph Minor Structure Theorem

This section introduces the graph minor structure theorem of Robertson and Seymour. This theorem shows that every graph in a proper minor-closed class can be constructed using four ingredients: graphs on surfaces, vortices, apex vertices, and clique-sums. We show that, with a restriction on the apex vertices, every graph that can be constructed using these ingredients has bounded layered treewidth, and thus admits layered separations of bounded width.

Let G be a graph embedded in a surface Σ . Let F be a facial cycle of G (thought of as a subgraph of G). An F -vortex is an F -decomposition $(B_x \subseteq V(H) : x \in V(F))$ of a graph H such that $V(G \cap H) = V(F)$ and $x \in B_x$ for each $x \in V(F)$. For $g, p, a \geq 0$ and $k \geq 1$, a graph G is (g, p, k, a) -almost-embeddable if for some set $A \subseteq V(G)$ with $|A| \leq a$, there are graphs

$G_0, G_1, \dots, G_s$ for some $s \in \{0, \dots, p\}$ such that:

- $G - A = G_0 \cup G_1 \cup \dots \cup G_s$,
- $G_1, \dots, G_s$ are pairwise vertex disjoint;
- G_0 is embeddable in a surface of Euler genus at most g ,
- there are s pairwise (vertex-)disjoint facial cycles $F_1, \dots, F_s$ of G_0 , and
- for $i \in \{1, \dots, s\}$, there is an F_i -vortex $(B_x \subseteq V(G_i) : x \in V(F_i))$ of G_i of width at most k .

The vertices in A are called *apex* vertices. They can be adjacent to any vertex in G .

A graph is *k-almost-embeddable* if it is (k, k, k, k) -almost-embeddable. The following graph minor structure theorem by Robertson and Seymour is at the heart of graph minor theory. In a tree decomposition $(B_x \subseteq V(G) : x \in V(T))$ of a graph G , the *torso* of a bag B_x is the subgraph obtained from $G[B_x]$ by adding all edges vw where $v, w \in B_x \cap B_y$ for some edge $xy \in E(T)$.

Theorem 16 (Robertson and Seymour [62]). *For every fixed graph H there is a constant $k = k(H)$ such that every H -minor-free graph is obtained by clique-sums of k -almost-embeddable graphs. Alternatively, every H -minor-free graph has a tree decomposition in which each torso is k -almost embeddable.*

This section explores which graphs described by the graph minor structure theorem admit layered separations of bounded width. As stated earlier, it is not the case that all such graphs admit layered separations of bounded width. For example, let G be the graph obtained from the $\sqrt{n} \times \sqrt{n}$ grid by adding one dominant vertex. Thus, G has diameter 2, contains no K_6 -minor, and has treewidth at least $\sqrt{n}$. By Lemma 5, if G admits layered separations of width ℓ , then $\ell \in \Omega(\sqrt{n})$.

We will show that the following restriction to the definition of almost-embeddable will lead to graph classes that admit layered separations of bounded width. A graph G is *strongly (g, p, k, a) -almost-embeddable* if it is (g, p, k, a) -almost-embeddable and there is no edge between an apex vertex and a vertex in $G_0 - (G_1 \cup \dots \cup G_s)$. That is, each apex vertex is only adjacent to other apex vertices or vertices in the vortices. A graph is *strongly k -almost-embeddable* if it is strongly (k, k, k, k) -almost-embeddable.

Theorem 17. *Every strongly (g, p, k, a) -almost-embeddable graph G is $(a + (k+1)(2g+2p+3))$ -good.*

Proof. We use the notation from the definition of strongly (g, p, k, a) -almost-embeddable. We may assume that G is connected, and except for $F_1, \dots, F_s$, each face of G_0 is a triangle, where

G_0 might contain parallel edges not bounding a single face. Let K be the given clique in G of size at most $a + (k + 1)(2g + 2p + 3)$ (in the definition of good).

Construct a layering $(V_0, V_1, \dots, V_t)$ of G as follows. Let $V_0 := K$ and let

$$V_1 := (N_G(K) \cup A \cup V(G_1 \cup \dots \cup G_s)) \setminus K.$$

For $i = 2, 3, \dots$, let V_i be the set of vertices of G that are not in $V_0 \cup \dots \cup V_{i-1}$ and are adjacent to some vertex in V_{i-1} . Thus $(V_0, V_1, \dots, V_t)$ is a layering of G (for some t).

Let $K' := (K \cap V(G_0) \setminus V(F_1 \cup \dots \cup F_s))$ be the part of K embedded in the surface and avoiding the vortices. If $K' \neq \emptyset$ then let r be one vertex in K' , otherwise r is undefined.

Let G'_0 be the triangulation obtained from G_0 as follows. For $i \in \{1, \dots, s\}$, add a new vertex r_i inside face F_i (corresponding to vortex G_i) and add an edge between r_i and each vertex of F_i . Let $n := |V(G'_0)|$.

We now construct a spanning forest T of G'_0 . Declare r (if defined) and $r_1, \dots, r_s$ to be the roots of T . For $i \in \{1, \dots, s\}$, make each vertex in $V(F_i)$ adjacent to r_i in T . By definition, these edges are in G'_0 . Now, make each vertex in $K' \setminus \{r\}$ adjacent to r in T . Since K' is a clique, these edges are in G'_0 . Every vertex v in $V(G'_0) \cap V_1$ that is not already in T is adjacent to K' ; make each such vertex v adjacent to a neighbour in K' in T . Every vertex in $V(G'_0) \cap V_1$ is now in T (either as a root or as a child or grandchild of a root). Now, for $i = 2, 3, \dots$, for each vertex v in $V(G'_0) \cap V_i$, choose a neighbour w of v in V_{i-1} , and add the edge vw to T . Now, T is a spanning forest of G'_0 with s or $s + 1$ connected components, and thus with $n - s$ or $n - s - 1$ edges.

Let D be the graph with vertex set $V(G'_0)$ where two vertices of D are adjacent if the corresponding faces share an edge in $G'_0 - E(T)$. Thus G'_0 has $3n + 3g - 6$ edges and $2n + 2g - 4$ faces. Hence $|V(D)| = 2n + 2g - 4$ and $|E(D)| = |E(G'_0)| - |E(T)| \leq (3n + 3g - 6) - (n - s - 1) = 2n + 3g + s - 5$. By the same argument used in the proof of Theorem 9, D is connected. Let T^* be a spanning tree of D . Let $X := E(D) \setminus E(T^*)$. Note that $|X| \leq (2n + 3g + s - 5) - (2n + 2g - 4 - 1) = g + s$.

For each vertex $x \in V(G'_0)$, let P_x be the path in T between x and the root of the connected component of T containing x . By construction, P_x includes at most one vertex in G'_0 in each layer V_i with $i \geq 1$. If P_x is in the component of T rooted at r , then let $P_x^+ := V(P_x) \setminus K$. Otherwise, P_x is in the component of T rooted at r_i for some $i \in \{1, \dots, s\}$. Then P_x contains exactly one vertex $v \in V(F_i \cap P_x)$. Let $P_x^+ := (V(P_x) \setminus \{r_i\}) \cup B_v$, where B_v is the bag indexed by v in the vortex G_i . Thus P_x^+ is a set of vertices in G with at most $k + 1$ vertices in each layer V_i with $i \geq 1$ (since $|B_v| \leq k + 1$). For each face $f = uvw$ of G'_0 , let

$$C_f := \cup \{P_x^+ \cup P_y^+ : xy \in X\} \cup P_u^+ \cup P_v^+ \cup P_w^+ \cup A \cup K.$$

Thus C_f contains at most $a + (k + 1)(2g + 2s + 3)$ vertices in each layer V_i (since $|K| \leq a + (k + 1)(2g + 2s + 3)$).

We now prove that $(C_f : f \in F(G'_0))$ is a T^* -decomposition of G . (This makes sense since $V(T^*) = F(G'_0)$.) First, we prove condition (1) in the definition of T^* -decomposition for each edge vw of G . If $v \in A \cup K$, then v is in every bag and w is in some bag (proved below), implying v and w are in a common bag. Now assume that $v \notin A \cup K$ and $w \notin A \cup K$ by symmetry. If $vw \in E(G_0)$, then $v, w \in C_f$ for each of the two faces f of G'_0 incident to vw . Otherwise $vw \in E(G_i)$ for some $i \in \{1, \dots, s\}$. Then $v, w \in B_x$ for some vertex $x \in V(F_i)$, implying that $v, w \in C_f$ for each face f of G'_0 incident to x (and there is such a face, for example, the one including r_i). This proves condition (1) in the definition of T^* -decomposition.

We now prove condition (2) in the definition of T^* -decomposition for each vertex v of G . We consider three cases: (a) v is in $A \cup K$, (b) v is in $G_0 - K - V(G_1 \cup \dots \cup G_s)$, or (c) v is in some vortex G_i .

(a) If $v \in A \cup K$, then v is in every bag, and condition (2) is satisfied for v .

(b) Say v is in $(G_0 - K) - V(G_1 \cup \dots \cup G_s)$. Let F' be the set of faces f of G'_0 such that v is in C_f . Each face incident to v is in F' , thus F' is non-empty. It now suffices to prove that the induced subgraph $T^*[F']$ is connected. Let T' be the subtree of T rooted at v . If some edge xy in X is a half-chord or chord of T' , then v is in $P_x \cup P_y$, implying that v is in every bag, and $T^*[F'] = T^*$ is connected. Now assume that no half-chord or chord of T' is in X . Then a face f of G'_0 is in F' if and only if f is incident with a vertex in T' ; that is, $F' = F(T')$. Let H be the graph defined in Lemma 8 with respect to T' . That is, H has vertex set F' and edge set the dual-chords and dual-half-chords of T' . By Lemma 8, H is connected. Since v is in $G_0 - K$, we have v is not a root of T . Let p be the parent of v in T . Each chord or half-chord of T' is an edge of $G - (E(T) \cup X)$, except for pv , which is a half-chord of T' (since $p \notin V(T')$). Let e be the edge of H dual to pv . By Lemma 8, $T^*[F'] = H - e$ is connected, as desired.

(c) Say v is in G_i for some $i \in \{1, \dots, s\}$. Let F' be the set of faces f of G'_0 such that v is in C_f . It suffices to prove that the induced subgraph $T^*[F']$ is connected and non-empty. Let $Z := \{z \in V(F_i) : v \in B_z\}$, where B_z is the bag of G_i corresponding to z . By the definition of a vortex, Z induces a connected non-empty subgraph of the cycle F_i . Say $Z = (z_1, z_2, \dots, z_q)$ ordered by F_i where $q \geq 1$. For $j \in \{1, \dots, q\}$, let T_j be the subtree of T rooted at z_j . Let F'_j be the set of faces of G'_0 incident to some vertex in T_j . By construction, $T^*[F'] = \bigcup_j T^*[F'_j]$. By the argument used in part (b) applied to z_j , $T^*[F'_j]$ is connected and non-empty. Since F'_j and F'_{j+1} have the face $r_i z_j z_{j+1}$ in common for $j \in \{1, \dots, q-1\}$, we have $T^*[F'] = \bigcup_j T^*[F'_j]$ is connected and non-empty, as desired.

Therefore $(C_f : f \in F(G'_0))$ is a T^* -decomposition of G , and it has layered width at most $a + (k+1)(2g+2s+3)$. $\square$

It is easily seen that every clique in a (g, p, k, a) -almost-embeddable graph has order at most $a + 2k + \lfloor \frac{1}{2}(7 + \sqrt{1 + 24g}) \rfloor$, which is at most $a + (k+1)(2g+2p+3)$ since $k \geq 1$ and $p \geq 0$. Thus Lemma 3, Lemma 15 and Theorem 17 together imply:

Theorem 18. *Every graph obtained by clique-sums of strongly (g, p, k, a) -almost-embeddable graphs is $a + (k + 1)(2g + 2p + 3)$ -good, has layered treewidth at most $a + (k + 1)(2g + 2p + 3)$, and admits layered separations of width $a + (k + 1)(2g + 2p + 3)$.*

Lemma 4 and Theorem 18 together imply:

Theorem 19. *Let G be a graph obtained by clique-sums of strongly k -almost-embeddable graphs. Then:*

- (a) *G is $(4k^2 + 8k + 3)$ -good,*
- (b) *G has layered treewidth at most $4k^2 + 8k + 3$,*
- (c) *G admits layered separations of width $4k^2 + 8k + 3$, and*
- (d) *if G has diameter d then G has treewidth less than $(4k^2 + 8k + 3)(d + 1)$.*

Theorem 19(d) improves upon a result by Grohe [40, Proposition 10] who proved an upper bound on the treewidth of $d \cdot f(k)$, where $f(k) \approx k^k$. Moreover, this result of Grohe [40] assumes there are no apex vertices. That is, it is for clique-sums of $(k, k, k, 0)$ -almost-embeddable graphs.

Recall that a graph H is *apex* if $H - v$ is planar for some vertex v of H . Dvorák and Thomas [34] proved a structure theorem for general H -minor-free graphs, which in the case of apex graphs H , says that H -minor-free graphs are obtained from clique-sums of strongly k -almost-embeddable graphs, for some $k = k(H)$; see [17] for related claims. Thus Theorem 19 implies:

Theorem 20. *For each fixed apex graph H there is a constant $\ell = \ell(H)$ such that every H -minor-free graph has layered treewidth at most ℓ and admits layered separations of width ℓ .*

We now characterise the minor-closed classes with bounded layered treewidth.

Theorem 21. *The following are equivalent for a proper minor-closed class of graphs $\mathcal{G}$:*

- (1) *every graph in $\mathcal{G}$ has bounded layered treewidth,*
- (2) *every graph in $\mathcal{G}$ admits layered separations of bounded width,*
- (3) *$\mathcal{G}$ has linear local treewidth,*
- (4) *$\mathcal{G}$ has bounded local treewidth,*
- (5) *$\mathcal{G}$ excludes a fixed apex graph as a minor,*
- (6) *every graph in $\mathcal{G}$ is obtained from clique-sums of strongly k -almost-embeddable graphs (for fixed k).*

Proof. Lemma 3 shows that (1) implies (2). Lemma 7 shows that (2) implies (3), which implies (4) by definition. Eppstein [36] proved that (4) and (5) are equivalent; see [15] for an alternative proof. As mentioned above, Dvorák and Thomas [34] proved that (5) implies (6). Theorem 19(b) proves that (6) implies (1). $\square$

Note that Demaine and Hajiaghayi [16] previously proved that (3) and (4) are equivalent.

6 Rich Decompositions and Shadow Complete Layerings

As observed in Section 5, it is not the case that graphs in every proper minor-closed class admit layered separations of bounded width. However, in this section we introduce some tools (namely, rich tree decompositions and shadow complete layerings) that enable our methods based on layered tree decompositions to be extended to conclude results about graphs in any proper minor-closed class. See Theorems 30 and 40 for two applications of the results in this section.

Consider a layering $(V_0, V_1, \dots, V_t)$ of a graph G . Let H be a connected component of $G[V_i \cup V_{i+1} \cup \dots \cup V_t]$, for some $i \in \{1, \dots, t\}$. The *shadow* of H is the set of vertices in V_{i-1} adjacent to H . The layering is *shadow complete* if every shadow is a clique. This concept was introduced by Kündgen and Pelsmayer [48] and implicitly by Dujmović et al. [26]. It is a key to the proof that graphs of bounded treewidth have bounded nonrepetitive chromatic number [48] and bounded track-number [26].

A tree decomposition $(B_x \subseteq V(G) : x \in V(T))$ of a graph G is *k-rich* if $B_x \cap B_y$ is a clique in G on at most k vertices, for each edge $xy \in E(T)$. The following lemma generalises a result by Kündgen and Pelsmayer [48], who proved it when each bag of the tree decomposition is a clique (that is, for chordal graphs). We allow bags to induce more general graphs, and in subsequent sections we apply this lemma with each bag inducing an ℓ -almost embeddable graph (Theorems 30 and 40).

For a subgraph H of a graph G , a tree decomposition $(C_y \subseteq V(H) : y \in V(F))$ of H is *contained* in a tree decomposition $(B_x \subseteq V(G) : x \in V(T))$ of G if for each bag C_y there is bag B_x such that $C_y \subseteq B_x$.

Lemma 22. *Let G be a graph with a k -rich tree decomposition $\mathcal{T}$ for some $k \geq 1$. Then G has a shadow complete layering $(V_0, V_1, \dots, V_t)$ such that every shadow has size at most k , and for each $i \in \{0, \dots, t\}$, the subgraph $G[V_i]$ has a $(k-1)$ -rich tree decomposition contained in $\mathcal{T}$.*

Proof. We may assume that G is connected with at least one edge. Say $\mathcal{T} = (B_x \subseteq V(G) : x \in V(T))$ is a k -rich tree decomposition of G . If $B_x \subseteq B_y$ for some edge $xy \in E(T)$, then contracting xy into y (and keeping bag B_y) gives a new k -rich tree decomposition of G . Moreover, if a tree decomposition of a subgraph of G is contained in the new tree decomposition of G , then

it is contained in the original. Thus, we may assume that $B_x \not\subseteq B_y$ and $B_y \not\subseteq B_x$ for each edge $xy \in V(T)$.

Let G' be the graph obtained from G by adding an edge between every pair of vertices in a common bag (if the edge does not already exist). Let r be a vertex of G . Let α be a node of T such that $r \in B_\alpha$. Root T at α . Now every non-root node of T has a parent node. Let $V_0 := \{r\}$. Let t be the eccentricity of r in G' . For $i \in \{1, \dots, t\}$, let V_i be the set of vertices of G at distance i from r in G' . Since G is connected, G' is connected. Thus $(V_0, V_1, \dots, V_t)$ is a layering of G' and also of G (since $G \subseteq G'$).

Since each bag B_x is a clique in G' , V_1 is the set of vertices of G in bags that contain r (not including r itself). More generally, V_i is the set of vertices of G in bags that intersects V_{i-1} but are not in $V_0 \cup \dots \cup V_{i-1}$.

Define $B'_\alpha := B_\alpha \setminus \{r\}$ and $B''_\alpha := \{r\}$. For a non-root node $x \in V(T)$ with parent node y , define $B'_x := B_x \setminus B_y$ and $B''_x := B_x \cap B_y$. Since $B_x \not\subseteq B_y$, we have $B'_x \neq \emptyset$. One should think that B'_x is the set of vertices that first appear in B_x when traversing down the tree decomposition from the root, while B''_x is the set of vertices in B_x that appear above x in the tree decomposition.

Consider a node x of T . Since B_x is a clique in G' , B_x is contained in at most two consecutive layers. Consider (not necessarily distinct) vertices $u, v \in B'_x$, which is not empty. Then the distance between u and r in G' equals the distance between v and r in G' . Thus B'_x is contained in one layer, say $V_{\ell(x)}$. Let w be the neighbour of v in some shortest path between B'_x and r in G' . Then w is in $B''_x \cap V_{\ell(x)-1}$. In conclusion, each bag B_x is contained in precisely two consecutive layers, $V_{\ell(x)-1} \cup V_{\ell(x)}$, such that $\emptyset \neq B'_x \subseteq V_{\ell(x)}$ and $B_x \cap V_{\ell(x)-1} \subseteq B''_x \neq \emptyset$. Also, observe that if y is an ancestor of x in T , then $\ell(y) \leq \ell(x)$. We call this property $(\star)$.

We now prove that $G[V_i]$ has the desired $(k-1)$ -rich tree decomposition. Since $G[V_0]$ has one vertex and no edges, this is trivial for $i = 0$. Now assume that $i \in \{1, \dots, t\}$.

Let T_i be the subgraph of T induced by the nodes x such that $\ell(x) \leq i$. By property $(\star)$, T_i is a (connected) subtree of T . We claim that $\mathcal{T}_i := (B_x \cap V_i : x \in V(T_i))$ is a T_i -decomposition of $G[V_i]$. First we prove that each vertex $v \in V_i$ is in some bag of $\mathcal{T}_i$. Let x be the node of T closest to α , such that $v \in B_x$. Then $v \in B'_x$ and $\ell(x) = i$. Hence v is in the bag $B_x \cap V_i$ of $\mathcal{T}_i$, as desired. Now we prove that for each edge $vw \in E(G[V_i])$, both v and w are in a common bag of $\mathcal{T}_i$. Let x be the node of T closest to α , such that $v \in B_x$. Let y be the node of T closest to α , such that $w \in B_y$. Since $vw \in E(G)$, there is a bag B_z containing both v and w , and z is a descendent of both x and y in T (by the definition of x and y). Thus, without loss of generality, x is on the $y\alpha$ -path in T . Thus $w \in B'_y$ and $y \in V(T_i)$. Moreover, v is also in B_y (since v and w are in a common bag of $\mathcal{T}$). Thus, v and w are in the bag $B_y \cap V_i$ of $\mathcal{T}_i$, as desired. Finally, we prove that for each vertex $v \in V_i$, the set of bags in $\mathcal{T}_i$ that contain v correspond to a (connected) subtree of T_i . By assumption, this property holds in T . Let X be the subtree of T whose corresponding bags in $\mathcal{T}$ contain v . Let x be the root of X . Then $v \in B'_x$ and $\ell(x) = i$. By property $(\star)$, we have $\ell(z) \in \{i, i+1\}$ for each node z in X . Moreover, again by property

($\star$), deleting from X the nodes z such that $\ell(z) = i + 1$ gives a connected subtree of X , which is precisely the subtree of T_i whose bags in $\mathcal{T}_i$ contain v . Hence $\mathcal{T}_i$ is a T_i -decomposition of $G[V_i]$. By definition, $\mathcal{T}_i$ is contained in $\mathcal{T}$.

We now prove that $\mathcal{T}_i$ is $(k - 1)$ -rich. Consider an edge $xy \in E(T_i)$. Without loss of generality, y is the parent of x in T_i . Our goal is to prove that $B_x \cap B_y \cap V_i = B''_x \cap V_i$ is a clique on at most $k - 1$ vertices. Certainly, it is a clique on at most k vertices, since $\mathcal{T}$ is k -rich. Now, $\ell(x) \leq i$ (since $x \in V(T_i)$). If $\ell(x) < i$ then $B_x \cap V_i = \emptyset$, and we are done. Now assume that $\ell(x) = i$. Thus $B'_x \subseteq V_i$ and $B'_x \neq \emptyset$. Let v be a vertex in B'_x . Let w be the neighbour of v on a shortest path in G' between v and r . Thus w is in $B''_x \cap V_{i-1}$. Thus $|B''_x \cap V_i| \leq k - 1$, as desired. Hence $\mathcal{T}_i$ is $(k - 1)$ -rich.

We now prove that $(V_0, V_1, \dots, V_t)$ is shadow complete. Let H be a connected component of $G[V_i \cup V_{i+1} \cup \dots \cup V_t]$ for some $i \in \{1, \dots, t\}$. Let X be the subgraph of T whose corresponding bags in $\mathcal{T}$ intersect $V(H)$. Since H is connected, X is indeed a connected subtree of T . Let x be the root of X . Consider a vertex w in the shadow of H . That is, $w \in V_{i-1}$ and w is adjacent to some vertex v in $V(H) \cap V_i$. Let y be the node closest to x in X , such that $v \in B_y$. Then $v \in B'_y$ and $w \in B''_y$. Thus $\ell(y) = i$. Consider a node z in X . Then $B_z \subseteq V_{\ell(z)-1} \cup V_{\ell(z)}$ and some vertex in B_z is in $V(H)$ and is thus in $V_i \cup V_{i+1} \cup \dots \cup V_t$. Thus $\ell(z) \geq i$. On the other hand, if in addition z is in the yx -path in X , then $\ell(z) \leq \ell(y) = i$ by property ($\star$), implying $\ell(z) = i$. Thus $w \in B''_z$ for each such node z . In particular, $w \in B''_x$. Since B''_x is a clique, the shadow of H is a clique. Hence $(V_0, V_1, \dots, V_t)$ is shadow complete. Moreover, since $|B''_x| \leq k$, the shadow of H has size at most k . $\square$

7 Track and Queue Layouts

The results of this section are expressed in terms of track layouts of graphs, which is a type of graph layout closely related to queue layouts and 3-dimensional grid drawings. A *vertex $|I|$ -colouring* of a graph G is a partition $\{V_i : i \in I\}$ of $V(G)$ such that for every edge $vw \in E(G)$, if $v \in V_i$ and $w \in V_j$ then $i \neq j$. The elements of the set I are *colours*, and each set V_i is a *colour class*. Suppose that $\preceq_i$ is a total order on each colour class V_i . Then each pair $(V_i, \preceq_i)$ is a *track*, and $\{(V_i, \preceq_i) : i \in I\}$ is an *$|I|$ -track assignment* of G .

An *X -crossing* in a track assignment consists of two edges vw and xy such that $v \prec_i x$ and $y \prec_j w$, for distinct colours i and j . A *t -track assignment* of G that has no X -crossings is called a *t -track layout* of G . The minimum t such that a graph G has t -track layout is called the *track-number* of G , denoted by $\text{tn}(G)$. Dujmović et al. [26] proved that

$$\text{qn}(G) \leq \text{tn}(G) - 1. \quad (1)$$

Conversely, Dujmović et al. [27] proved that $\text{tn}(G) \leq f(\text{qn}(G))$ for some function f . In this sense, queue-number and track-number are tied.

As described in Section 1.2, Dujmović [22] recently showed that layered separators can be used to construct queue layouts. In fact, the construction produces a track layout, which with (1) gives the desired bound for queue layouts.

Lemma 23 ([22]). *If a graph G admits layered separations of width ℓ then*

$$\text{qn}(G) < \text{tn}(G) \leq 3\ell(\lceil \log_{3/2} n \rceil + 1) .$$

Recall the following result discussed in Section 1.1.

Lemma 24 ([23, 49]). *Every planar graph admits layered separations of width 2.*

Lemmas 23 and 24 imply the following result of Dujmović [22].

Theorem 25 ([22]). *Every n -vertex planar graph G satisfies*

$$\text{qn}(G) < \text{tn}(G) \leq 6\lceil \log_{3/2} n \rceil + 6 .$$

This bound on $\text{qn}(G)$ was improved to $\text{qn}(G) \leq 4\lceil \log_{3/2} n \rceil$ by Fabrizio Frati [personal communication, 2013].

Now consider queue and track layouts of graphs with Euler genus g . Theorem 10 and Lemma 23 imply that $\text{qn}(G) < \text{tn}(G) \in \mathcal{O}(g \log n)$. This bound can be improved to $\mathcal{O}(g + \log n)$ as follows. A straightforward extension of the proof of Lemma 23 gives the following result (see Appendix C for a proof):

Lemma 26. *Let $\mathcal{T}$ be a tree decomposition of a graph G , such that there is a set $Q \subseteq V(G)$ with at most ℓ_1 vertices in each layer of some layering of G , and $\mathcal{T}$ restricted to $G - Q$ has layered width at most ℓ_2 with respect to the same layering. Then*

$$\text{qn}(G) < \text{tn}(G) \leq 3\ell_1 + 3\ell_2(1 + \log_{3/2} n) .$$

Theorem 13 and Lemma 26 with $\ell_1 = 2g$ and $\ell_2 = 3$ imply the following generalisation of the above results.

Theorem 27. *For every n -vertex graph with Euler genus g ,*

$$\text{qn}(G) < \text{tn}(G) \leq 6g + 9(1 + \log_{3/2} n) .$$

Theorem 20 and Lemma 23 imply the following further generalisation.

Theorem 28. *For each fixed apex graph H , for every n -vertex H -minor-free graph G ,*

$$\text{qn}(G) < \text{tn}(G) \leq \mathcal{O}(\log n) .$$

We now extend this result to arbitrary proper minor-closed classes. Dujmović et al. [26] implicitly proved that if a graph G has a shadow complete layering such that each layer induces a graph with track-number at most c and each shadow has size at most s , then G has track-number at most $3c^{s+1}$ (see Appendix D). Iterating this result gives the next lemma.

Lemma 29 ([26]). *For some number c , let $\mathcal{G}_0$ be a class of graphs with track-number at most c . For $k \geq 1$, let $\mathcal{G}_k$ be a class of graphs that have a shadow complete layering such that each shadow has size at most k , and each layer induces a graph in $\mathcal{G}_{k-1}$. Then every graph in $\mathcal{G}_k$ has track-number and queue-number at most $3^{(k+2)!} c^{(k+1)!}$.*

Theorem 30. *For every fixed graph H , every H -minor-free n -vertex graph has track-number and queue-number at most $\log^{\mathcal{O}(1)} n$.*

Proof. Let G_0 be an H -minor-free graph on n vertices. By Theorem 16, there are constants $k \geq 1$ and $\ell \geq 1$ depending only on H , such that G_0 is a spanning subgraph of a graph G that has a k -rich tree decomposition $\mathcal{T}$ such that each bag induces an ℓ -almost embeddable subgraph of G . Note that G_0 , G , $\mathcal{T}$ and n remain fixed throughout the proof.

For $j \in \{0, \dots, k\}$, let $\mathcal{G}_j$ be the set of induced subgraphs of G that have a j -rich tree decomposition contained in $\mathcal{T}$. Note that G itself is in $\mathcal{G}_k$.

Consider a graph $G' \in \mathcal{G}_0$. Then G' is the union of disjoint subgraphs of G , each of which is ℓ -almost embeddable. To layout one ℓ -almost embeddable graph, put each of the at most ℓ apex vertices on its own track, and layout the remaining graph with $3(4\ell^2 + 8\ell + 3)(\lceil \log_{3/2} n \rceil + 1)$ tracks by Theorem 19 and Lemma 23. (Here we do not use the clique-sums in Theorem 19.) Of course, the track-number of a graph is the maximum track-number of its connected components. Thus G' has track-number at most $\ell + 3(4\ell^2 + 8\ell + 3)(\lceil \log_{3/2} n \rceil + 1)$.

Consider $G' \in \mathcal{G}_j$ for some $j \in \{1, \dots, k\}$. Thus G' is an induced subgraph of G with a j -rich tree decomposition contained in $\mathcal{T}$. By Lemma 22, G' has a shadow complete layering $(V_0, \dots, V_t)$ such that every shadow has size at most j , and for each layer V_i , the induced subgraph $G'[V_i]$ has a $(j-1)$ -rich tree decomposition $\mathcal{T}_i$ contained in $\mathcal{T}$. Thus $G'[V_i]$ is in $\mathcal{G}_{j-1}$.

By Lemma 29 with $c = \ell + 3(4\ell^2 + 8\ell + 3)(\lceil \log_{3/2} n \rceil + 1)$, our graph G has track-number at most $3^{(k+2)!} (\ell + 3(4\ell^2 + 8\ell + 3)(\lceil \log_{3/2} n \rceil + 1))^{(k+1)!}$, which is in $\log^{\mathcal{O}(1)} n$ since k and ℓ are constants (depending only on H).

The claimed bound on queue-number follows from (1). □

8 3-Dimensional Graph Drawing

This section presents our results for 3-dimensional graph drawings, which are based on the following connection between track layouts and 3-dimensional graph drawings.

Lemma 31 ([26, 30]). *If a c -colourable n -vertex graph G has a t -track layout, then G has 3-dimensional grid drawings with $\mathcal{O}(t^2n)$ volume and with $\mathcal{O}(c^7tn)$ volume.*

Every graph with Euler genus g is $\mathcal{O}(\sqrt{g})$ -colourable [47]. Thus Theorem 27 and Lemma 31 imply:

Theorem 32. *Every n -vertex graph with Euler genus g has a 3-dimensional grid drawing with volume $\mathcal{O}(g^{7/2}(g + \log n)n)$.*

For fixed H , every H -minor-free graph is $\mathcal{O}(1)$ -colourable. Thus, Theorem 28 and Lemma 31 imply:

Theorem 33. *For each fixed apex graph H , every n -vertex H -minor-free graph has a 3-dimensional grid drawing with volume $\mathcal{O}(n \log n)$.*

Lemma 31 and Theorem 30 extend this theorem to arbitrary proper minor-closed classes:

Theorem 34. *For each fixed graph H , every H -minor-free n -vertex graph has a 3-dimensional grid drawing with volume $n \log^{\mathcal{O}(1)} n$.*

The best previous upper bound on the volume of 3-dimensional grid drawings of graphs with bounded Euler genus or H -minor-free graphs was $\mathcal{O}(n^{3/2})$ [30].

9 Nonrepetitive Colourings

This section proves our results for nonrepetitive colourings. Recall the following two results by Dujmović et al. [23] discussed in Section 1.3. (Theorem 36 is implied by Lemmas 24 and 35.)

Lemma 35 ([23]). *If an n -vertex graph G admits layered separations of width ℓ then*

$$\pi(G) \leq 4\ell(1 + \log_{3/2} n) .$$

Theorem 36 ([23]). *For every n -vertex planar graph G ,*

$$\pi(G) \leq 8(1 + \log_{3/2} n) .$$

Now consider nonrepetitive colourings of graphs G with Euler genus g . Theorem 10 and Lemma 35 imply that $\pi(G) \leq \mathcal{O}(g \log n)$. This bound can be improved to $\mathcal{O}(g + \log n)$ as follows. A straightforward extension of the proof of Lemma 35 gives the following result (see Appendix C for a proof):

Lemma 37. *Let $\mathcal{T}$ be a tree decomposition of a graph G , such that there is a set $Q \subseteq V(G)$ with at most ℓ_1 vertices in each layer of some layering of G , and $\mathcal{T}$ restricted to $G - Q$ has layered width at most ℓ_2 with respect to the same layering. Then*

$$\pi(G) \leq 4\ell_1 + 4\ell_2(1 + \log_{3/2} n) .$$

Theorem 13 and Lemma 37 with $\ell_1 = 2g$ and $\ell_2 = 3$ imply the following generalisation of the above results.

Theorem 38. *For every n -vertex graph with Euler genus g ,*

$$\pi(G) \leq 8g + 12(1 + \log_{3/2} n) .$$

To generalise Theorem 38, we employ a result by Kündgen and Pelsmajer [48]. They proved that if a graph G has a shadow complete layering such that the graph induced by each layer is nonrepetitively c -colourable, then G is nonrepetitively $4c$ -colourable [48, Theorem 6]. Iterating this result gives the next lemma.

Lemma 39 ([48]). *For some number c , let $\mathcal{G}_0$ be a class of graphs with nonrepetitive chromatic number at most c . For $k \geq 1$, let $\mathcal{G}_k$ be a class of graphs that have a shadow complete layering such that each layer induces a graph in $\mathcal{G}_{k-1}$. Then every graph in $\mathcal{G}_k$ has nonrepetitive chromatic number at most $c4^k$.*

The next theorem is our main result about nonrepetitive colourings.

Theorem 40. *For every fixed graph H , every H -minor-free n -vertex graph is nonrepetitively $\mathcal{O}(\log n)$ -colourable.*

Proof. Let G_0 be an H -minor-free graph on n vertices. By Theorem 16, there are constants $k \geq 1$ and $\ell \geq 1$ depending only on H , such that G_0 is a spanning subgraph of a graph G that has a k -rich tree decomposition $\mathcal{T}$ such that each bag induces an ℓ -almost embeddable subgraph of G . Note that G_0 , G , $\mathcal{T}$ and n remain fixed throughout the proof.

For $j \in \{0, \dots, k\}$, let $\mathcal{G}_j$ be the set of induced subgraphs of G that have a j -rich tree decomposition contained in $\mathcal{T}$. Note that G itself is in $\mathcal{G}_k$.

Consider a graph $G' \in \mathcal{G}_0$. Then G' is the union of disjoint subgraphs of G , each of which is ℓ -almost embeddable. To nonrepetitively colour one ℓ -almost embeddable graph, give each of the at most ℓ apex vertices its own colour and colour the remainder with $4(4\ell^2 + 8\ell + 3)(1 + \log_{3/2} n)$ colours by Theorem 19 and Lemma 35. (Here we do not use the clique-sums in Theorem 19.) Of course, colours can be reused on distinct connected components. Thus G' is nonrepetitively $(\ell + 4(4\ell^2 + 8\ell + 3)(1 + \log_{3/2} n))$ -colourable.

Consider some $G' \in \mathcal{G}_j$ for some $j \in \{1, \dots, k\}$. Thus G' is an induced subgraph of G with a j -rich tree decomposition contained in $\mathcal{T}$. By Lemma 22, G' has a shadow complete layering

$(V_0, \dots, V_t)$ such that for each layer V_i , the induced subgraph $G'[V_i]$ has a $(j-1)$ -rich tree decomposition $\mathcal{T}_i$ contained in $\mathcal{T}$. Thus $G'[V_i]$ is in $\mathcal{G}_{j-1}$.

By Lemma 39 with $c = \ell + 4(4\ell^2 + 8\ell + 3)(1 + \log_{3/2} n)$, our graph G is nonrepetitively $4^k(\ell + 4(4\ell^2 + 8\ell + 3)(1 + \log_{3/2} n))$ -colourable, as is G_0 , since G_0 is a subgraph of G . $\square$

10 Reflections

1. The notion of local treewidth has been very successful in the field of approximation algorithms and bidimensionality [4, 14, 16, 40]. Given that layered tree decompositions can be thought of as a global structure for graphs of bounded local treewidth, it would be interesting to see if layered treewidth has applications to approximation algorithms and bidimensionality.

2. What can be said about the layered treewidth of various non-minor-closed graph classes?

3. The similarity between queue/track layouts and nonrepetitive colourings is remarkable given how different the definitions seem at first glance. Both parameters have bounded expansion [52] and admit very similar properties with respect to subdivisions [31, 52]. Many proof techniques work for both queue/track layouts and nonrepetitive colourings, in particular layered separations and shadow-complete layerings. One exception is that graphs of bounded maximum degree have bounded nonrepetitive chromatic number [2, 24, 41, 44], whereas graphs of bounded maximum degree have unbounded track- and queue-number [70]. It would be interesting to prove a more direct relationship. Do graphs of bounded track/queue-number have bounded nonrepetitive chromatic number? More specifically, do 1-queue graphs have bounded nonrepetitive chromatic number? And do 3-track graphs have bounded nonrepetitive chromatic number?

4. Finally, we mention the work of Shahrokhi [63] who introduced a definition equivalent to layered treewidth. (We became aware of reference [63] when it was posted on the arXiv in 2015.) Shahrokhi [63] was motivated by questions completely different from those in the present paper. In our language, he proved that for every graph G with layered treewidth k , there is a graph G_1 with clique cut width at most $2k - 1$ and a chordal graph G_2 such that $G = G_1 \cap G_2$. Shahrokhi [63] then proved that every planar graph G has layered treewidth at most 4, implying that there is a graph G_1 with clique cut width at most 7 and a chordal graph G_2 such that $G = G_1 \cap G_2$. Theorem 9 with $g = 0$ improves these bounds from 4 to 3 and thus from 7 to 5. All our other results about layered treewidth can be applied in this domain as well.

Acknowledgements

Thanks to Zdenek Dvorák, Gwenaél Joret, Sergey Norin, Bruce Reed and Paul Seymour for helpful discussions. This research was partially completed at Bellairs Research Institute in Barbados.

References

- [1] LYUDMIL G. ALEKSANDROV AND HRISTO N. DJIDJEV. Linear algorithms for partitioning embedded graphs of bounded genus. *SIAM J. Discrete Math.*, 9(1):129–150, 1996. doi: [10.1137/S0895480194272183](https://doi.org/10.1137/S0895480194272183). MR: [1375420](#).
- [2] NOGA ALON, JAROSŁAW GRZYCZUK, MARIUSZ HAŁUSZCZAK, AND OLIVER RIORDAN. Non-repetitive colorings of graphs. *Random Structures Algorithms*, 21(3–4):336–346, 2002. doi: [10.1002/rsa.10057](https://doi.org/10.1002/rsa.10057). MR: [1945373](#).
- [3] NOGA ALON, PAUL D. SEYMOUR, AND ROBIN THOMAS. A separator theorem for nonplanar graphs. *J. Amer. Math. Soc.*, 3(4):801–808, 1990. doi: [10.2307/1990903](https://doi.org/10.2307/1990903). MR: [1065053](#), Zbl: [0747.05051](#).
- [4] BRENDA S. BAKER. Approximation algorithms for NP-complete problems on planar graphs. *J. Assoc. Comput. Mach.*, 41(1):153–180, 1994. doi: [10.1145/174644.174650](https://doi.org/10.1145/174644.174650). MR: [1369197](#).
- [5] JÁNOS BARÁT AND PÉTER P. VARIÚ. On square-free vertex colorings of graphs. *Studia Sci. Math. Hungar.*, 44(3):411–422, 2007. doi: [10.1556/SScMath.2007.1029](https://doi.org/10.1556/SScMath.2007.1029). MR: [2361685](#).
- [6] NORMAN BIGGS. Spanning trees of dual graphs. *J. Combinatorial Theory Ser. B*, 11:127–131, 1971. MR: [0284369](#).
- [7] ROBIN BLANKENSHIP. *Book Embeddings of Graphs*. Ph.D. thesis, Department of Mathematics, Louisiana State University, U.S.A., 2003. http://etd.lsu.edu/docs/available/etd-0709103-163907/unrestricted/Blankenship_dis.pdf.
- [8] PROSENJIT BOSE, JUREK CZYZOWICZ, PAT MORIN, AND DAVID R. WOOD. The maximum number of edges in a three-dimensional grid-drawing. *J. Graph Algorithms Appl.*, 8(1):21–26, 2004. doi: [10.7155/jgaa.00079](https://doi.org/10.7155/jgaa.00079).
- [9] JEAN BOURGAIN. Expanders and dimensional expansion. *C. R. Math. Acad. Sci. Paris*, 347(7–8):357–362, 2009. doi: [10.1016/j.crma.2009.02.009](https://doi.org/10.1016/j.crma.2009.02.009). MR: [2537230](#).
- [10] JEAN BOURGAIN AND AMIR YEHUDAYOFF. Expansion in $SL_2(\mathbb{R})$ and monotone expansion. *Geometric and Functional Analysis*, 23(1):1–41, 2013. doi: [10.1007/s00039-012-0200-9](https://doi.org/10.1007/s00039-012-0200-9). MR: [3037896](#).
- [11] BOŠTJAN BREŠAR, JAROSŁAW GRZYCZUK, SANDI KLAVŽAR, STANISŁAW NIWCZYK, AND IZTOK PETERIN. Nonrepetitive colorings of trees. *Discrete Math.*, 307(2):163–172, 2007. doi: [10.1016/j.disc.2006.06.017](https://doi.org/10.1016/j.disc.2006.06.017). MR: [2285186](#).
- [12] PANAGIOTIS CHEILARIS, ERNST SPECKER, AND STATHIS ZACHOS. Neochromatica. *Comment. Math. Univ. Carolin.*, 51(3):469–480, 2010. <http://www.dml.cz/dmlcz/140723>. MR: [2741880](#).
- [13] ROBERT F. COHEN, PETER EADES, TAO LIN, AND FRANK RUSKEY. Three-dimensional graph drawing. *Algorithmica*, 17(2):199–208, 1996. doi: [10.1007/BF02522826](https://doi.org/10.1007/BF02522826).

- [14] ERIK D. DEMAINE, FEDOR V. FOMIN, MOHAMMADTAGHI HAJIAGHAYI, AND DIMITRIOS M. THILIKOS. Bidimensional parameters and local treewidth. *SIAM J. Discrete Math.*, 18(3):501–511, 2004/05. doi: [10.1137/S0895480103433410](https://doi.org/10.1137/S0895480103433410). MR: [2134412](#).
- [15] ERIK D. DEMAINE AND MOHAMMADTAGHI HAJIAGHAYI. Diameter and treewidth in minor-closed graph families, revisited. *Algorithmica*, 40(3):211–215, 2004. doi: [10.1007/s00453-004-1106-1](https://doi.org/10.1007/s00453-004-1106-1). MR: [2080518](#).
- [16] ERIK D. DEMAINE AND MOHAMMADTAGHI HAJIAGHAYI. Equivalence of local treewidth and linear local treewidth and its algorithmic applications. In *Proc. 15th Annual ACM-SIAM Symposium on Discrete Algorithms (SODA '04)*, pp. 840–849. SIAM, 2004. <http://dl.acm.org/citation.cfm?id=982792.982919>.
- [17] ERIK D. DEMAINE, MOHAMMADTAGHI HAJIAGHAYI, AND KEN-ICHI KAWARABAYASHI. Approximation algorithms via structural results for apex-minor-free graphs. In *36th International Colloquium on Automata, Languages and Programming (ICALP '09)*, pp. 316–327. 2009. doi: [10.1007/978-3-642-02927-1_27](https://doi.org/10.1007/978-3-642-02927-1_27).
- [18] GIUSEPPE DI BATTISTA, FABRIZIO FRATI, AND JANOS PÁCH. On the queue number of planar graphs. *SIAM J. Comput.*, 42(6):2243–2285, 2013. doi: [10.1137/130908051](https://doi.org/10.1137/130908051). MR: [3141759](#).
- [19] EMILIO DI GIACOMO, GIUSEPPE LIOTTA, HENK MEIJER, AND STEPHEN K. WISMATH. Volume requirements of 3D upward drawings. *Discrete Mathematics*, 309(7):1824–1837, 2009. doi: [10.1016/j.disc.2007.12.061](https://doi.org/10.1016/j.disc.2007.12.061). MR: [2509356](#).
- [20] REINHARD DIESTEL AND DANIELA KÜHN. Graph minor hierarchies. *Discrete Appl. Math.*, 145(2):167–182, 2005. doi: [10.1016/j.dam.2004.01.010](https://doi.org/10.1016/j.dam.2004.01.010).
- [21] HRISTO N. DJIDIEV. On the constants of separator theorems. *C. R. Acad. Bulgare Sci.*, 40(10):31–34, 1987. MR: [940049](#).
- [22] VIDA DUJMOVIĆ. Graph layouts via layered separators. *J. Combin. Theory Series B.*, 110:79–89, 2015. doi: [10.1016/j.jctb.2014.07.005](https://doi.org/10.1016/j.jctb.2014.07.005). arXiv: [1302.0304](https://arxiv.org/abs/1302.0304). MR: [3279388](#).
- [23] VIDA DUJMOVIĆ, FABRIZIO FRATI, GWENAËL JORET, AND DAVID R. WOOD. Nonrepetitive colourings of planar graphs with $O(\log n)$ colours. *Electronic J. Combinatorics*, 20(1):P51, 2013. <http://www.combinatorics.org/ojs/index.php/eljc/article/view/v20i1p51>.
- [24] VIDA DUJMOVIĆ, GWENAËL JORET, JAKUB KOZIK, AND DAVID R. WOOD. Nonrepetitive colouring via entropy compression. 2012. To appear in *Combinatorica*, arXiv: [1112.5524](https://arxiv.org/abs/1112.5524).
- [25] VIDA DUJMOVIĆ, PAT MORIN, AND ADAM SHEFFER. Crossings in grid drawings. *Electron. J. Combin.*, 21(1):#P1.41, 2014. <http://www.combinatorics.org/ojs/index.php/eljc/article/view/v21i1p41>.
- [26] VIDA DUJMOVIĆ, PAT MORIN, AND DAVID R. WOOD. Layout of graphs with bounded tree-width. *SIAM J. Comput.*, 34(3):553–579, 2005. doi: [10.1137/S0097539702416141](https://doi.org/10.1137/S0097539702416141). MR: [2137079](#), Zbl: [1069.05055](#).

- [27] VIDA DUJMOVIĆ, ATTILA PÓR, AND DAVID R. WOOD. Track layouts of graphs. *Discrete Math. Theor. Comput. Sci.*, 6(2):497–522, 2004. <http://www.dmtcs.org/dmtcs-ojs/index.php/dmtcs/article/view/204>. MR: 2180055.
- [28] VIDA DUJMOVIĆ, ANASTASIOS SIDIROPOULOS, AND DAVID R. WOOD. 3-monotone expanders. 2015. arXiv: [1501.05020](https://arxiv.org/abs/1501.05020).
- [29] VIDA DUJMOVIĆ AND DAVID R. WOOD. On linear layouts of graphs. *Discrete Math. Theor. Comput. Sci.*, 6(2):339–358, 2004. <http://www.dmtcs.org/dmtcs-ojs/index.php/dmtcs/article/view/193>. MR: 2081479.
- [30] VIDA DUJMOVIĆ AND DAVID R. WOOD. Three-dimensional grid drawings with sub-quadratic volume. In JÁNOS PACH, ed., *Towards a Theory of Geometric Graphs*, vol. 342 of *Contemporary Mathematics*, pp. 55–66. Amer. Math. Soc., 2004. MR: 2065252.
- [31] VIDA DUJMOVIĆ AND DAVID R. WOOD. Stacks, queues and tracks: Layouts of graph subdivisions. *Discrete Math. Theor. Comput. Sci.*, 7:155–202, 2005. <http://www.dmtcs.org/dmtcs-ojs/index.php/dmtcs/article/view/67>. MR: 2164064.
- [32] VIDA DUJMOVIĆ AND DAVID R. WOOD. Upward three-dimensional grid drawings of graphs. *Order*, 23(1):1–20, 2006. doi: [10.1007/s11083-006-9028-y](https://doi.org/10.1007/s11083-006-9028-y). MR: 2258457.
- [33] ZEEV DVIR AND AVI WIGDERSON. Monotone expanders: Constructions and applications. *Theory of Computing*, 6(12):291–308, 2010. doi: [10.4086/toc.2010.v006a012](https://doi.org/10.4086/toc.2010.v006a012). MR: 2770077.
- [34] ZDENEK DVORÁK AND ROBIN THOMAS. List-coloring apex-minor-free graphs. 2013. arXiv: [1401.1399](https://arxiv.org/abs/1401.1399).
- [35] DAVID EPPSTEIN. Subgraph isomorphism in planar graphs and related problems. *J. Graph Algorithms Appl.*, 3(3):1–27, 1999. doi: [10.7155/jgaa.00014](https://doi.org/10.7155/jgaa.00014). MR: 1750082.
- [36] DAVID EPPSTEIN. Diameter and treewidth in minor-closed graph families. *Algorithmica*, 27(3–4):275–291, 2000. doi: [10.1007/s004530010020](https://doi.org/10.1007/s004530010020). MR: 1759751.
- [37] DAVID EPPSTEIN. Dynamic generators of topologically embedded graphs. In *Proc. 14th Annual ACM-SIAM Symposium on Discrete Algorithms, SODA '03*, pp. 599–608. SIAM, 2003. <http://dl.acm.org/citation.cfm?id=644108.644208>.
- [38] STEFAN FELSNER, GIUSEPPE LIOTTA, AND STEPHEN K. WISMATH. Straight-line drawings on restricted integer grids in two and three dimensions. In PETRA MUTZEL, MICHAEL JÜNGER, AND SEBASTIAN LEIPERT, eds., *Proc. 9th International Symp. on Graph Drawing (GD '01)*, vol. 2265 of *Lecture Notes in Comput. Sci.*, pp. 328–342. Springer, 2002.
- [39] JOHN R. GILBERT, JOAN P. HUTCHINSON, AND ROBERT E. TARJAN. A separator theorem for graphs of bounded genus. *J. Algorithms*, 5(3):391–407, 1984. doi: [10.1016/0196-6774\(84\)90019-1](https://doi.org/10.1016/0196-6774(84)90019-1).
- [40] MARTIN GROHE. Local tree-width, excluded minors, and approximation algorithms. *Combinatorica*, 23(4):613–632, 2003. doi: [10.1007/s00493-003-0037-9](https://doi.org/10.1007/s00493-003-0037-9). MR: 2046826.

- [41] JAROS LAW GRZYTCZUK. Nonrepetitive colorings of graphs—a survey. *Int. J. Math. Math. Sci.*, 74639, 2007. doi: [10.1155/2007/74639](https://doi.org/10.1155/2007/74639). MR: [2272338](#).
- [42] JAROS LAW GRZYTCZUK. Thue type problems for graphs, points, and numbers. *Discrete Math.*, 308(19):4419–4429, 2008. doi: [10.1016/j.disc.2007.08.039](https://doi.org/10.1016/j.disc.2007.08.039). MR: [2433769](#).
- [43] RUDOLF HALIN. S -functions for graphs. *J. Geometry*, 8(1-2):171–186, 1976. doi: [10.1007/BF01917434](https://doi.org/10.1007/BF01917434). MR: [0444522](#).
- [44] JOCHEN HARANTA AND STANISLAV JENDROL'. Nonrepetitive vertex colorings of graphs. *Discrete Math.*, 312(2):374–380, 2012. doi: [10.1016/j.disc.2011.09.027](https://doi.org/10.1016/j.disc.2011.09.027). MR: [2852595](#).
- [45] LENWOOD S. HEATH, F. THOMSON LEIGHTON, AND ARNOLD L. ROSENBERG. Comparing queues and stacks as mechanisms for laying out graphs. *SIAM J. Discrete Math.*, 5(3):398–412, 1992. doi: [10.1137/0405031](https://doi.org/10.1137/0405031). MR: [1172748](#).
- [46] LENWOOD S. HEATH AND ARNOLD L. ROSENBERG. Laying out graphs using queues. *SIAM J. Comput.*, 21(5):927–958, 1992. doi: [10.1137/0221055](https://doi.org/10.1137/0221055). MR: [1181408](#).
- [47] PERCY J. HEAWOOD. Map colour theorem. *Quart. J. Pure Appl. Math.*, 24:332–338, 1890. doi: [10.1112/plms/s2-51.3.161](https://doi.org/10.1112/plms/s2-51.3.161).
- [48] ANDRE KÜNDGEN AND MICHAEL J. PELSMAJER. Nonrepetitive colorings of graphs of bounded tree-width. *Discrete Math.*, 308(19):4473–4478, 2008. doi: [10.1016/j.disc.2007.08.043](https://doi.org/10.1016/j.disc.2007.08.043). MR: [2433774](#).
- [49] RICHARD J. LIPTON AND ROBERT E. TARIAN. A separator theorem for planar graphs. *SIAM J. Appl. Math.*, 36(2):177–189, 1979. doi: [10.1137/0136016](https://doi.org/10.1137/0136016). MR: [0524495](#).
- [50] SETH M. MALITZ. Genus g graphs have pagewidth $O(\sqrt{g})$. *J. Algorithms*, 17(1):85–109, 1994. doi: [10.1006/jagm.1994.1028](https://doi.org/10.1006/jagm.1994.1028). MR: [1279270](#).
- [51] BOJAN MOHAR AND CARSTEN THOMASSEN. *Graphs on surfaces*. Johns Hopkins University Press, Baltimore, U.S.A., 2001.
- [52] JAROSLAV NEŠETŘIL, PATRICE OSSONA DE MENDEZ, AND DAVID R. WOOD. Characterisations and examples of graph classes with bounded expansion. *European J. Combinatorics*, 33(3):350–373, 2011. doi: [10.1016/j.ejc.2011.09.008](https://doi.org/10.1016/j.ejc.2011.09.008). MR: [2864421](#).
- [53] L. TAYLOR OLLMANN. On the book thicknesses of various graphs. In FREDERICK HOFFMAN, ROY B. LEVOW, AND ROBERT S. D. THOMAS, eds., *Proc. 4th Southeastern Conference on Combinatorics, Graph Theory and Computing*, vol. VIII of *Congr. Numer.*, p. 459. Utilitas Math., 1973.
- [54] JÁNOS PACH, TORSTEN THIELE, AND GÉZA TÓTH. Three-dimensional grid drawings of graphs. In BERNARD CHAZELLE, JACOB E. GOODMAN, AND RICHARD POLLACK, eds., *Advances in discrete and computational geometry*, vol. 223 of *Contemporary Mathematics*, pp. 251–255. Amer. Math. Soc., 1999.

- [55] SRIRAM V. PEMMARAJU. *Exploring the Powers of Stacks and Queues via Graph Layouts*. Ph.D. thesis, Virginia Polytechnic Institute and State University, U.S.A., 1992.
- [56] BRUCE A. REED. Tree width and tangles: a new connectivity measure and some applications. In *Surveys in combinatorics*, vol. 241 of *London Math. Soc. Lecture Note Ser.*, pp. 87–162. Cambridge Univ. Press, 1997. doi: [10.1017/CBO9780511662119.006](https://doi.org/10.1017/CBO9780511662119.006).
- [57] S. RENGARAJAN AND C. E. VENI MADHAVAN. Stack and queue number of 2-trees. In DING-ZHU DU AND MING LI, eds., *Proc. 1st Annual International Conf. on Computing and Combinatorics (COCOON '95)*, vol. 959 of *Lecture Notes in Comput. Sci.*, pp. 203–212. Springer, 1995. doi: [10.1007/BFb0030834](https://doi.org/10.1007/BFb0030834).
- [58] BRUCE RICHTER AND HERBERT SHANK. The cycle space of an embedded graph. *J. Graph Theory*, 8(3):365–369, 1984. doi: [10.1002/jgt.3190080304](https://doi.org/10.1002/jgt.3190080304). MR: [754916](https://www.jstor.org/stable/754916).
- [59] NEIL ROBERTSON AND PAUL D. SEYMOUR. Graph minors. III. Planar tree-width. *J. Combin. Theory Ser. B*, 36(1):49–64, 1984. doi: [10.1016/0095-8956\(84\)90013-3](https://doi.org/10.1016/0095-8956(84)90013-3). MR: [742386](https://www.jstor.org/stable/742386).
- [60] NEIL ROBERTSON AND PAUL D. SEYMOUR. Graph minors. II. Algorithmic aspects of tree-width. *J. Algorithms*, 7(3):309–322, 1986. doi: [10.1016/0196-6774\(86\)90023-4](https://doi.org/10.1016/0196-6774(86)90023-4).
- [61] NEIL ROBERTSON AND PAUL D. SEYMOUR. Excluding a graph with one crossing. In *Graph Structure Theory. Proc. of AMS-IMS-SIAM Joint Summer Research Conference on Graph Minors*, vol. 147 of *Contemp. Math.*, pp. 669–675. Amer. Math. Soc., 1993. doi: [10.1090/conm/147/01206](https://doi.org/10.1090/conm/147/01206). MR: [1224738](https://www.jstor.org/stable/1224738).
- [62] NEIL ROBERTSON AND PAUL D. SEYMOUR. Graph minors. XVI. Excluding a non-planar graph. *J. Combin. Theory Ser. B*, 89(1):43–76, 2003. doi: [10.1016/S0095-8956\(03\)00042-X](https://doi.org/10.1016/S0095-8956(03)00042-X).
- [63] FARHAD SHAHROKHI. New representation results for planar graphs. In *29th European Workshop on Computational Geometry (EuroCG 2013)*, pp. 177–180. 2013. arXiv: [1502.06175](https://arxiv.org/abs/1502.06175).
- [64] MARTIN ŠKOVIERA. Spanning subgraphs of embedded graphs. *Czechoslovak Math. J.*, 42(117)(2):235–239, 1992. MR: [1179495](https://www.jstor.org/stable/1179495).
- [65] AXEL THUE. Über unendliche Zeichenreihen. *Norske Vid. Selsk. Skr. I. Mat. Nat. Kl. Christiania*, 7:1–22, 1906.
- [66] K. TRUEMPER. *Matroid Decomposition*. Academic Press, Boston, 1992. MR: [1170126](https://www.jstor.org/stable/1170126).
- [67] KARL GEORG CHRISTIAN VON STAUDT. *Geometrie der Lage*. Verlag von Bauer and Rappe 25. Julius Merz, Nürnberg, 1847.
- [68] KLAUS WAGNER. Über eine Eigenschaft der ebene Komplexe. *Math. Ann.*, 114:570–590, 1937. doi: [10.1007/BF01594196](https://doi.org/10.1007/BF01594196). MR: [1513158](https://www.jstor.org/stable/1513158). Zbl: [0017.19005](https://zbmath.org/journals/0017.19005).

- [69] DAVID R. WOOD. Queue layouts of graph products and powers. *Discrete Math. Theor. Comput. Sci.*, 7(1):255–268, 2005. <http://www.dmtcs.org/dmtcs-ojs/index.php/dmtcs/article/view/71>.
- [70] DAVID R. WOOD. Bounded-degree graphs have arbitrarily large queue-number. *Discrete Math. Theor. Comput. Sci.*, 10(1):27–34, 2008. <http://www.dmtcs.org/dmtcs-ojs/index.php/dmtcs/article/viewArticle/541>. MR: 2369152.
- [71] DAVID R. WOOD. Clique minors in cartesian products of graphs. *New York J. Math.*, 17:627–682, 2011. <http://nyjm.albany.edu/j/2011/17-28.html>.
- [72] DAVID R. WOOD AND JAN ARNE TELLE. Planar decompositions and the crossing number of graphs with an excluded minor. *New York J. Math.*, 13:117–146, 2007. <http://nyjm.albany.edu/j/2007/13-8.html>.
- [73] MIHALIS YANNAKAKIS. Embedding planar graphs in four pages. *J. Comput. System Sci.*, 38(1):36–67, 1989. doi: [10.1016/0022-0000\(89\)90032-9](https://doi.org/10.1016/0022-0000(89)90032-9). MR: 0990049.

A Separators in a Tree Decomposition

For completeness we include the following proof:

Proof of Lemma 1. Let $(B_x \subseteq V(G) : x \in V(T))$ be a T -decomposition of a graph G , for some tree T . Let $S \subseteq V(G)$. Our goal is to prove that there is a bag B such that each connected component of $G - B$ contains at most $\frac{1}{2}|S|$ vertices in S .

Consider an edge xy of T . Let $T(x, y)$ and $T(y, x)$ be the components of $T - xy$ that respectively contain x and y . Let $G(x, y)$ be the subgraph of $G - B_y$ induced by the union of the bags associated with nodes in $T(x, y)$. Define $G(y, x)$ analogously. Observe that each vertex of G is in exactly one of $G(x, y)$, $G(y, x)$ and $B_x \cap B_y$. Without loss of generality, $G(x, y)$ contains at most $\frac{1}{2}|S|$ vertices in S . Orient xy from x to y .

Since T is a tree, there is a node y of T , such that each edge of T incident to y is oriented towards y . That is, for each neighbour x of y in T , the subgraph $G(x, y)$ contains at most $\frac{1}{2}|S|$ vertices in S . Thus, each connected component of $G - B_y$ contains at most $\frac{1}{2}|S|$ vertices in S , and B_y is the desired bag. This proves the first claim.

We now show that G has a separation (G_1, G_2) with $V(G_1 \cap G_2) = B$ and both $G_1 - V(G_2)$ and $G_2 - V(G_1)$ contain at most $\frac{2}{3}|S|$ vertices in S . Partition the components of $G - B_y$ into two parts P and Q to minimise the maximum number of vertices in S in one of the parts. Let p and q be the number of vertices in S in P and Q respectively, where $p \geq q$. Suppose that $p > \frac{2}{3}|S|$. Thus $q < \frac{1}{3}|S| < \frac{1}{2}p$. Each component in P contains at most $\frac{1}{2}|S|$ vertices in S . Thus

P has at least two components and therefore has a component C that contains at least 1 and at most $\frac{1}{2}p$ vertices in S . Let P', Q' be the partition obtained by moving C from P to Q . Thus P' contains less vertices in S than P . And Q' contains at most $q + \frac{1}{2}p$ vertices in S , which is less than p . This contradicts the choice of P, Q . Hence $q \leq p \leq \frac{2}{3}|S|$. Thus G contains the desired separation. $\square$

B Layered Treewidth to Treewidth

The following result is due to Sergey Norin [personal communication, 2014].

Theorem 41. *Every n -vertex graph G with layered treewidth k has treewidth at most $2\sqrt{kn}$.*

Proof. Let $V_1, V_2, \dots, V_t$ be the layering in a tree decomposition of G with layered width k . Let $p := \lceil \sqrt{n/k} \rceil$. For $j \in \{1, \dots, p\}$, let $W_j := V_j \cup V_{p+j} \cup V_{2p+j} \cup \dots$. Thus $W_1, W_2, \dots, W_p$ is a partition of $V(G)$. For some $j \in \{1, \dots, p\}$ we have $|W_j| \leq \frac{n}{p} \leq \sqrt{kn}$. Each connected component of $G - W_j$ is contained within $p - 1$ consecutive layers, and therefore has treewidth at most $k(p - 1) - 1 \leq \sqrt{kn}$. Hence $G - W_j$ has treewidth at most $\sqrt{kn}$. Adding W_j to every bag gives a tree decomposition of G with width at most $\sqrt{kn} + |W_j| \leq 2\sqrt{kn}$. $\square$

C Recursive Separators

Here we prove Lemmas 26 and 37. The method, which is based on recursive application of layered separations, is a straightforward generalisation of the method of Dujmović et al. [23] for nonrepetitive colouring and of Dujmović [22] for track layouts. Both lemmas have the same starting assumptions: Let $V_1, V_2, \dots, V_p$ be a layering of a graph G . Let $\mathcal{T}$ be a tree decomposition of G , such that there is a set $Q \subseteq V(G)$ with at most ℓ_1 vertices in each layer V_i , and $\mathcal{T}$ restricted to $G - Q$ has layered width at most ℓ_2 with respect to $V_1, V_2, \dots, V_p$.

For each vertex $v \in Q$, let $\text{depth}(v) := 0$. For $i \in \{1, \dots, p\}$, injectively label the vertices in $V_i \cap Q$ by $1, 2, \dots, \ell_1$. Let $\text{label}(v)$ be the label assigned to each vertex $v \in V_i \cap Q$. By assumption, $G - Q$ has layered treewidth at most ℓ_2 and thus admits layered separations of width ℓ_2 by Lemma 3. Now run the following recursive algorithm $\text{COMPUTE}(V(G) \setminus Q, 1)$.

COMPUTE (input S and d , where $S \subseteq V(G) \setminus Q$ and $d \in \mathbb{Z}^+$)

1. If $S = \emptyset$ then exit.
2. Let (G_1, G_2) be a separation of $G - Q$ such that each layer V_i contains at most ℓ_2 vertices in $V(G_1 \cap G_2) \cap S$, and both $V(G_1) - V(G_2)$ and $V(G_2) - V(G_1)$ contain at most $\frac{2}{3}|S|$ vertices in S .
3. Let $\text{depth}(v) := d$ for each vertex $v \in V(G_1 \cap G_2) \cap S$.
4. For $i \in \{1, \dots, p\}$, injectively label the vertices in $V_i \cap V(G_1 \cap G_2) \cap S$ by $1, 2, \dots, \ell_2$. Let $\text{label}(v)$ be the label assigned to each vertex $v \in V_i \cap V(G_1 \cap G_2) \cap S$.
5. COMPUTE($(V(G_1) - V(G_2)) \cap S, d + 1$)
6. COMPUTE($(V(G_2) - V(G_1)) \cap S, d + 1$)

The recursive application of COMPUTE determines a rooted binary tree T , where each node of T corresponds to one call to COMPUTE. Associate each vertex whose depth and label is computed in a particular call to COMPUTE with the corresponding node of T . (Observe that the depth and label of each vertex is determined exactly once.) Note that the maximum depth is at most $1 + \log_{3/2} n$.

Proof of Lemma 26. Our goal is to prove that $\text{tn}(G) \leq 3\ell_1 + 3\ell_2(1 + \log_{3/2} n)$. The tracks are indexed by triples of integers as follows. Colour each vertex v by $(\text{col}(v), \text{depth}(v), \text{label}(v))$, where $\text{col}(v) := i \bmod 3$ if $v \in V_i$, and depth and label are computed above. This defines a track assignment for G . We now order each track. Consider two vertices $v \in V_i$ and $w \in V_j$ on the same track; that is, $(\text{col}(v), \text{depth}(v), \text{label}(v)) = (\text{col}(w), \text{depth}(w), \text{label}(w))$. If $i < j$ then place $v \prec w$ in the track. If $j < i$ then place $w \prec v$ in the track. Now assume that $i = j$. If v and w are associated with the same node of T , since $i = j$ we have $\text{label}(v) \neq \text{label}(w)$, which is a contradiction. Now assume v and w are associated with distinct nodes of T with least common ancestor α . Say S was the input set corresponding to α , and (G_1, G_2) was the corresponding separation of $G - Q$. Without loss of generality, $v \in (V(G_1) - V(G_2)) \cap S$ and $w \in (V(G_2) - V(G_1)) \cap S$. Place $v \prec w$ in the track. It is easily seen that each track is totally ordered by $\preceq$.

We now show there is no X-crossing. Suppose that edges vw and xy form an X-crossing, where $(\text{col}(v), \text{depth}(v), \text{label}(v)) = (\text{col}(x), \text{depth}(x), \text{label}(x))$ and $(\text{col}(w), \text{depth}(w), \text{label}(w)) = (\text{col}(y), \text{depth}(y), \text{label}(y))$ and $v \prec x$ and $y \prec w$. Say $v \in V_a$ and $w \in V_b$ and $x \in V_c$ and $y \in V_d$. Since vw and xy are edges, $|a - b| \leq 1$ and $|c - d| \leq 1$. Since $\text{col}(v) = \text{col}(x)$ and $\text{col}(w) = \text{col}(y)$ we have $a \equiv c \pmod{3}$ and $b \equiv d \pmod{3}$. Since $v \prec x$ and $y \prec w$ we have $a \leq c$ and $d \leq b$. If $a < c$ then $a + 3 \leq c \leq d + 1 \leq b + 1 \leq a + 2$, which is a contradiction. Similarly, if $d < b$ then $d + 3 \leq b \leq a + 1 \leq c + 1 \leq d + 2$, which is a contradiction. Now assume

that $a = c$ and $d = b$. Without loss of generality, $\text{depth}(v) = \text{depth}(x) \leq \text{depth}(w) = \text{depth}(y)$. Since $\text{label}(v) = \text{label}(x)$ and $v \neq x$, we have v and x are associated with distinct nodes of T . Let α be the least common ancestor of these nodes of T . Say S was the input set corresponding to α , and (G_1, G_2) was the corresponding separation of $G - Q$. Since $v \prec x$ we have $v \in (V(G_1) - V(G_2)) \cap S$ and $x \in (V(G_2) - V(G_1)) \cap S$. Since $\text{depth}(v) \leq \text{depth}(w)$ and vw is an edge, $w \in (V(G_1) - V(G_2)) \cap S$. Similarly, since $\text{depth}(x) \leq \text{depth}(y)$ and xy is an edge, $y \in (V(G_2) - V(G_1)) \cap S$. Therefore the algorithm places $x \prec y$ on their track, which is a contradiction. Hence no two edges form an X-crossing. The number of tracks is at most $3\ell_1 + 3\ell_2(1 + \log_{3/2} n)$. $\square$

Proof of Lemma 37. Our goal is to prove that $\pi(G) \leq 4\ell_1 + 4\ell_2(1 + \log_{3/2} n)$. Kündgen and Pelsmayer [48] proved that for every layering of a graph G , there is a (not necessarily proper) 4-colouring of G such that for every repetitively coloured path $(v_1, v_2, \dots, v_{2t})$, the subpaths $(v_1, v_2, \dots, v_t)$ and $(v_{t+1}, v_{t+2}, \dots, v_{2t})$ have the same layer pattern (that is, for $i \in \{1, \dots, t\}$, vertices v_i and v_{t+i} are in the same layer). Let col be a such a 4-colouring. Now colour each vertex v by $(\text{col}(v), \text{depth}(v), \text{label}(v))$, where depth and label are computed above. Suppose on the contrary that $(v_1, v_2, \dots, v_{2t})$ is a repetitively coloured path in G . Then $(v_1, v_2, \dots, v_t)$ and $(v_{t+1}, v_{t+2}, \dots, v_{2t})$ have the same layer pattern. In addition, $\text{depth}(v_i) = \text{depth}(v_{t+i})$ and $\text{label}(v_i) = \text{label}(v_{t+i})$ for all $i \in [1, t]$. Let v_i and v_{t+i} be vertices in this path with minimum depth. Since v_i and v_{t+i} are in the same layer and have the same label, these two vertices were not labelled at the same step of the algorithm. Let x and y be the two nodes of T respectively associated with v_i and v_{t+i} . Let z be the least common ancestor of x and y in T . Say node z corresponds to call $\text{COMPUTE}(B, d)$. Thus v_i and v_{t+i} are in B (since if a vertex v is in B in the call to COMPUTE associated with some node q of T , then v is in B in the call to COMPUTE associated with each ancestor of q in T). Let (G_1, G_2) be the separation in $\text{COMPUTE}(B, d)$. Since $\text{depth}(v_i) = \text{depth}(v_{t+i}) > d$, neither v_i nor v_{t+i} are in $V(G_1 \cap G_2)$. Since z is the least common ancestor of x and y , without loss of generality, $v_i \in V(G_1) - V(G_2)$ and $v_{t+i} \in V(G_2) - V(G_1)$. Thus some vertex v_j in the subpath $(v_{i+1}, v_{i+2}, \dots, v_{t+i-1})$ is in $V(G_1 \cap G_2)$. If $v_j \in B$ then $\text{depth}(v_j) = d$. If $v_j \notin B$ then $\text{depth}(v_j) < d$. In both cases, $\text{depth}(v_j) < \text{depth}(v_i) = \text{depth}(v_{t+i})$, which contradicts the choice of v_i and v_{t+i} . Hence there is no repetitively coloured path in G . There are $4\ell_1$ colours at depth 0 and $4\ell_2$ colours at every other depth. Since the maximum depth is at most $1 + \log_{3/2} n$, the number of colours is at most $4\ell_1 + 4\ell_2(1 + \log_{3/2} n)$. $\square$

Note that in both Lemmas 26 and 37 we may replace $\log_{3/2} n$ by $\log_2 n$ by using separators (and the first part of Lemma 1) instead of separations (as in the second part of Lemma 1).

D Track Layout Construction

Here we sketch a proof of a result used in Section 7 that is implicit in the work of Dujmović et al. [26].

Lemma 42 (Dujmović et al. [26]). *If a graph G has a shadow complete layering $V_1, \dots, V_t$ such that each layer induces a graph with track-number at most c and each shadow has size at most s , then G has track-number at most $3c^{s+1}$.*

Proof Sketch. Let T be the graph obtained from G by contracting each connected component of each subgraph $G[V_i]$ into a single node. For each node x of T , let H_x be the corresponding connected component. Let V'_i be the vertices of T arising from V_i . Thus $V'_1, \dots, V'_t$ is a layering of T . For each node $y \in V'_i$ where $i \in \{1, \dots, t\}$, let C_y be the set of neighbours of H_y in V_{i-1} . We may assume that $C_y \neq \emptyset$. Since the given layering is shadow complete, C_y is a clique, called the *parent clique* of y . Now C_y is contained in a single connected component H_x of $G[V_{i-1}]$, for some node $x \in V'_{i-1}$. Call x the *parent node* and H_x the *parent component* of y . This shows that each node in V'_i has exactly one neighbour in V'_{i-1} , which implies that T is a forest. As illustrated in Figure 2, T has a 3-track layout T_0, T_1, T_2 .




Figure 2: A 3-track layout of T .

By assumption, for each node x of T , there is a c -track layout of H_x . For a clique C of H_x of size at most s , define the *signature* of C to be the set of (at most s) tracks that contain C . Since there is no X-crossing, the set of cliques of H_x with the same signature can be linearly ordered $C_1 \prec \dots \prec C_p$, such that if v and w are vertices in the same track, and in distinct cliques C_i and C_j with $i < j$, then $v \prec w$ in that track. Call this a *clique ordering*.

Replace each track T_j of T by c sub-tracks, and replace each node $x \in T_j$ by the c -track layout of H_x . This defines a $3c$ track assignment for G . Clearly an edge in some H_x crosses no other edge. Two edges between a parent component H_x and the same child component H_y do not form an X-crossing, since the endpoints in H_x of such edges form a clique (the parent clique of y), and therefore are in distinct tracks. The only possible X-crossing is between edges ab and cd , where a and c are in some parent component H_x , and b and d are in distinct child components H_y and H_z , respectively.

To solve this problem, when determining the 3-track layout of T , the child nodes of each node x are ordered in their track so that $y \prec z$ whenever the parent cliques C_y and C_z have the same signature, and $C_y \prec C_z$ in the clique ordering. Then group the child nodes of x according to the signatures of their parent cliques, and for each signature σ , use a distinct set of c tracks for the child components whose parent cliques have signature σ . Now the ordering of the child components with the same signature agrees with the clique ordering of their parent cliques, and therefore agrees with the ordering of any neighbours in the parent component. It follows that there is no X-crossing. The number of tracks is at most $3c$ times the number of signatures, which is at most $\sum_{i=1}^s \binom{c}{i} \leq c^s$. In total there are at most $3c \cdot c^s$ tracks. $\square$

This proof makes no effort to reduce the number of tracks. Various tricks due to Dujmović et al. [26] and Di Giacomo et al. [19] make a modest improvement.