

# On the Metric Dimension of Cartesian Products of Graphs<sup>\*</sup>

José Cáceres<sup>a</sup> Carmen Hernando<sup>b</sup> Merce Mora<sup>b</sup>  
Ignacio M. Pelayo<sup>b</sup> María L. Puertas<sup>a</sup> Carlos Seara<sup>b</sup>  
David R. Wood<sup>b</sup>

<sup>a</sup>*Departamento de Estadística y Matemática Aplicada  
Universidad de Almería, Almería, Spain  
{jcaceres,mpuertas}@ual.es*

<sup>b</sup>*Departament de Matemàtica Aplicada II, Universitat Politècnica de Catalunya,  
Barcelona, Spain  
{carmen.hernando,merce.mora,ignacio.m.pelayo,carlos.seara,david.wood}@upc.edu*

---

## Abstract

A set  $S$  of vertices in a graph  $G$  *resolves*  $G$  if every vertex is uniquely determined by its vector of distances to the vertices in  $S$ . The *metric dimension* of  $G$  is the minimum cardinality of a resolving set of  $G$ . This paper studies the metric dimension of cartesian products  $G \square H$ . We prove that the metric dimension of  $G \square G$  is tied in a strong sense to the minimum order of a so-called doubly resolving set in  $G$ . Using bounds on the order of doubly resolving sets, we establish bounds on  $G \square H$  for many examples of  $G$  and  $H$ . One of our main results is a family of graphs  $G$  with bounded metric dimension for which the metric dimension of  $G \square G$  is unbounded.

*Key words:* graph, distance, resolving set, metric dimension, metric basis, cartesian product, Hamming graph, Mastermind, coin weighing

---

## 1 Introduction

This paper undertakes a general study of the metric dimension of cartesian products of graphs. All the graphs considered are finite, undirected, simple,

---

<sup>\*</sup> Research supported by projects MCYT-FEDER-BFM2003-00368, Gen-Cat-2001SGR00224, MCYT-HU2002-0010, MTM-2004-07891-C02-01, MEC-SB2003-0270

and connected. (The results can easily be generalised to disconnected graphs—we omit the details.) The vertex set and edge set of a graph  $G$  are denoted by  $V(G)$  and  $E(G)$ . The distance between vertices  $v, w \in V(G)$  is denoted by  $d_G(v, w)$ , or  $d(v, w)$  if the graph  $G$  is clear from the context. A vertex  $x \in V(G)$  *resolves* a pair of vertices  $v, w \in V(G)$  if  $d(v, x) \neq d(w, x)$ . A set of vertices  $S \subseteq V(G)$  *resolves*  $G$ , and  $S$  is a *resolving set* of  $G$ , if every pair of distinct vertices of  $G$  are resolved by some vertex in  $S$ . A resolving set  $S$  of  $G$  with the minimum cardinality is a *metric basis* of  $G$ , and  $|S|$  is the *metric dimension* of  $G$ , denoted by  $\beta(G)$ .

The *cartesian product* of graphs  $G$  and  $H$ , denoted by  $G \square H$ , is the graph with vertex set  $V(G) \times V(H) := \{(a, v) : a \in V(G), v \in V(H)\}$ , where  $(a, v)$  is adjacent to  $(b, w)$  whenever  $a = b$  and  $\{v, w\} \in E(H)$ , or  $v = w$  and  $\{a, b\} \in E(G)$ . Where there is no confusion the vertex  $(a, v)$  of  $G \square H$  will be written  $av$ . Observe that if  $G$  and  $H$  are connected, then  $G \square H$  is connected. In particular, for all vertices  $av, bw$  of  $G \square H$  we have  $d(av, bw) = d_G(a, b) + d_H(v, w)$ . Assuming isomorphic graphs are equal, the cartesian product is associative, and  $G_1 \square G_2 \square \dots \square G_d$  is well-defined.

Resolving sets in general graphs were first defined by Harary and Melter [19] and Slater [33], although as we shall see, resolving sets in hypercubes were studied earlier under the guise of a coin weighing problem [1, 5, 6, 13, 14, 18, 20, 23, 24, 25, 26, 27, 35]. Resolving sets have since been widely investigated [4, 7, 8, 9, 10, 12, 21, 28, 29, 30, 31, 32, 34, 36, 37, 38], and arise in many diverse areas including network discovery and verification [2], robot navigation [21, 32], connected joins in graphs [31], and strategies for the Mastermind game [3, 11, 15, 16, 17, 20].

Part of our motivation for studying the metric dimension of cartesian products is that in two of the above-mentioned applications, namely Mastermind and coin weighing, the graphs that arise are in fact cartesian products. These connections are explained in Sections 2 and 6 respectively.

The main contributions of this paper are based on the notion of doubly resolving sets, which are introduced in Section 4. We prove that the minimum order of a doubly resolving set in a graph  $G$  is tied in a strong sense to  $\beta(G \square G)$ . Thus doubly resolving sets are essential in the study of metric dimension of cartesian products. We then give a number of examples of bounds on the metric dimension of cartesian products through doubly resolving sets. In particular, Sections 5, 6, 7, 8, and 9 respectively study complete graphs, Hamming graphs, paths and grids, cycles, and trees. One of our main results here is a family of (highly connected) graphs with bounded metric dimension for which the metric dimension of the cartesian product is unbounded.

## 2 Coin Weighing and Hypercubes

The *hypercube*  $Q_n$  is the graph whose vertices are the  $n$ -dimensional binary vectors, where two vertices are adjacent if they differ in exactly one coordinate. It is well known that

$$Q_n = \underbrace{K_2 \square K_2 \square \dots \square K_2}_n.$$

It is easily seen that  $\beta(Q_n) \leq n$  (see Equation (11)). The first case when this bound is not tight is  $n = 5$ . A laborious calculation verifies that  $Q_5$  is resolved by the 4-vertex set  $\{(0, 0, 0, 0, 0), (0, 0, 0, 1, 1), (0, 0, 1, 0, 1), (0, 1, 0, 0, 1)\}$ . We have determined  $\beta(Q_n)$  for small values of  $n$  by computer search.

$n$	2	3	4	5	6	7	8	10	15
$\beta(Q_n)$	2	3	4	4	5	6	6	$\leq 7$	$\leq 10$

The asymptotic value of  $\beta(Q_n)$  turns out to be related to the following coin weighing problem first posed by Söderberg and Shapiro [35]. (See [18] for a survey on various coin weighing problems.) Given  $n$  coins, each with one of two distinct weights, determine the weight of each coin with the minimum number of weighings. We are interested in the static variant of this problem, where the choice of sets of coins to be weighed is determined in advance. Weighing a set  $S$  of coins determines how many light (and heavy) coins are in  $S$ , and no further information. It follows that the minimum number of weighings differs from  $\beta(Q_n)$  by at most one [20, 31]. A lower bound on the number of weighings by Erdős and Rényi [13] and an upper bound by Lindström [23] imply that

$$\lim_{n \rightarrow \infty} \beta(Q_n) \cdot \frac{\log n}{n} = 2,$$

where, as always in this paper, logarithms are binary. Note that Lindström's proof is constructive. He gives a scheme of  $2^k - 1$  weighings that suffice for  $k \cdot 2^{k-1}$  coins.

## 3 Projections

Let  $S$  be a set of vertices in the cartesian product  $G \square H$  of graphs  $G$  and  $H$ . The *projection* of  $S$  onto  $G$  is the set of vertices  $a \in V(G)$  for which there exists a vertex  $av \in S$ . Similarly, the *projection* of  $S$  onto  $H$  is the set of vertices  $v \in V(H)$  for which there exists a vertex  $av \in S$ . A *column* of  $G \square H$  is a set of vertices  $\{av : v \in V(H)\}$  for some vertex  $a \in V(G)$ , and a *row* of  $G \square H$  is a set of vertices  $\{av : a \in V(G)\}$  for some vertex  $v \in V(H)$ . Observe that each row induces a copy of  $G$ , and each column induces a copy of  $H$ . This

terminology is consistent with a representation of  $G \square H$  by the points of the  $|V(G)| \times |V(H)|$  grid.

**Lemma 1** *Let  $S \subseteq V(G \square H)$  for graphs  $G$  and  $H$ . Then every pair of vertices in a fixed row of  $G \square H$  are resolved by  $S$  if and only if the projection of  $S$  onto  $G$  resolves  $G$ . Similarly, every pair of vertices in a fixed column of  $G \square H$  are resolved by  $S$  if and only if the projection of  $S$  onto  $H$  resolves  $H$ .*

**Proof.** Consider two vertices  $av$  and  $aw$  in a common column. For every other vertex  $bx$  of  $G \square H$ , we have  $d(av, bx) - d(aw, bx) = d_H(v, x) - d_H(w, x)$ . Thus  $d(av, bx) \neq d(aw, bx)$  if and only if  $d_H(v, x) \neq d_H(w, x)$ . That is,  $av$  and  $aw$  are resolved by  $bx$  if and only if  $v$  and  $w$  are resolved by  $x$  in  $H$ . Hence  $av$  and  $aw$  are resolved by  $S$  if and only if  $v$  and  $w$  are resolved by the projection of  $S$  onto  $H$ . We have the analogous result for the projection onto  $G$  by symmetry.  $\square$

**Corollary 2** *For all graphs  $G$  and  $H$ , and for every resolving set  $S$  of  $G \square H$ , the projection of  $S$  onto  $G$  resolves  $G$ , and the projection of  $S$  onto  $H$  resolves  $H$ . In particular,  $\beta(G \square H) \geq \max\{\beta(G), \beta(H)\}$ .  $\square$*

## 4 Doubly Resolving Sets

Many of the results that follow are based on the following definitions. Let  $G \neq K_1$  be a graph. Two vertices  $v, w$  of  $G$  are *doubly resolved* by  $x, y \in V(G)$  if

$$d(v, x) - d(w, x) \neq d(v, y) - d(w, y).$$

A set of vertices  $S \subseteq V(G)$  *doubly resolves*  $G$ , and  $S$  is a *doubly resolving set*, if every pair of distinct vertices  $v, w \in V(G)$  are doubly resolved by two vertices in  $S$ . Every graph with at least two vertices has a doubly resolving set. Let  $\psi(G)$  denote the minimum cardinality of a doubly resolving set of a graph  $G \neq K_1$ . Note that if  $x, y$  doubly resolves  $v, w$  then  $d(v, x) - d(w, x) \neq 0$  or  $d(v, y) - d(w, y) \neq 0$ , and at least one of  $x$  and  $y$  (singly) resolves  $v, w$ . Thus a doubly resolving set is also a resolving set, and

$$\beta(G) \leq \psi(G).$$

Our interest in doubly resolving sets is based on the following upper bound.

**Theorem 3** *For all graphs  $G$  and  $H \neq K_1$ ,*

$$\beta(G \square H) \leq \beta(G) + \psi(H) - 1.$$

**Proof.** Let  $S$  be a metric basis of  $G$ . Let  $T$  be a doubly resolving set of  $H$  with  $|T| = \psi(H)$ . Fix vertices  $s \in S$  and  $t \in T$ . Let

$$X := \{sv : v \in T\} \cup \{at : a \in S\}.$$

Observe that  $|X| = |S| + |T| - 1$ . To prove that  $X$  resolves  $G \square H$ , consider two vertices  $av$  and  $bw$  of  $G \square H$ . By Lemma 1, if  $a = b$  then  $av$  and  $bw$  are resolved since the projection of  $X$  onto  $H$  is  $T$ . Similarly, if  $v = w$  then  $av$  and  $bw$  are resolved since the projection of  $X$  onto  $G$  is  $S$ . Now assume that  $a \neq b$  and  $v \neq w$ . Since  $T$  is doubly resolving for  $H$ , there are two vertices  $x, y \in T$  such that

$$d_H(v, x) - d_H(w, x) \neq d_H(v, y) - d_H(w, y).$$

Thus for at least one of  $x$  and  $y$ , say  $x$ , we have

$$d_H(v, x) - d_H(w, x) \neq d_G(b, s) - d_G(a, s).$$

Hence

$$d(av, sx) = d_G(a, s) + d_H(v, x) \neq d_G(b, s) + d_H(w, x) = d(bw, sx).$$

That is,  $sx \in X$  resolves  $av$  and  $bw$ .  $\square$

The relationship between resolving sets of cartesian products and doubly resolving sets is strengthened by the following lower bound.

**Lemma 4** *Suppose that  $S$  resolves  $G \square G$  for some graph  $G$ . Let  $A$  and  $B$  be the two projections of  $S$  onto  $G$ . Then  $A \cup B$  doubly resolves  $G$ . In particular,*

$$\beta(G \square G) \geq \frac{1}{2}\psi(G).$$

**Proof.** For any two vertices  $v, w \in V(G)$ , there is a vertex  $pq \in S$  that resolves  $vw, wv$ . That is,  $d(vw, pq) \neq d(wv, pq)$ . Thus  $d(v, p) + d(w, q) \neq d(w, p) + d(v, q)$ , which implies  $d(v, p) - d(w, p) \neq d(v, q) - d(w, q)$ . Thus  $p, q$  doubly resolves  $v, w$  in  $G$ . Now  $p \in A$  and  $q \in B$ . Hence  $A \cup B$  doubly resolves  $G$ . If  $S$  is a metric basis of  $G$ , then  $\psi(G) \leq |A \cup B| \leq |A| + |B| \leq 2|S| = 2 \cdot \beta(G \square G)$ .  $\square$

Observe that Theorem 3 and Lemma 4 prove that  $\beta(G \square G)$  is always within a constant factor of  $\psi(G)$ . In particular,

$$\frac{1}{2}\psi(G) \leq \beta(G \square G) \leq \psi(G) + \beta(G) - 1 \leq 2\psi(G) - 1. \quad (1)$$

Thus doubly resolving sets are essential in the study of the metric dimension of cartesian products.

A natural candidate for a resolving set of  $G \square G$  is  $S \times S$  for a well chosen set  $S \subseteq V(G)$ . It follows from Lemma 4 and the proof technique employed in Theorem 3 that  $S \times S$  resolves  $G \square G$  if and only if  $S$  doubly resolves  $G$ .

## 5 Complete Graphs

Let  $K_n$  denote the complete graph on  $n \geq 1$  vertices. Chartrand *et al.* [8] proved that for every  $n$ -vertex graph  $G$ ,

$$\beta(G) = n - 1 \iff G = K_n. \quad (2)$$

**Lemma 5** *For all  $n \geq 2$  we have  $\psi(K_n) = \max\{n - 1, 2\}$ .*

**Proof.** Since  $\psi(G) \geq 2$  for every graph  $G \neq K_1$ , we have  $\psi(K_2) = 2$ . Now suppose that  $n \geq 3$ . Thus  $\psi(K_n) \geq n - 1$  by (2). Let  $S$  be a set of all but one vertex in  $K_n$ . To prove that  $S$  doubly resolves  $K_n$ , consider two vertices  $v, w$  of  $K_n$ . Without loss of generality  $v \in S$ . Since  $n \geq 3$  there is a vertex  $y \in S \setminus \{v\}$ . Then  $d(v, v) - d(w, v) = 0 - 1 = -1$  and  $d(v, y) - d(w, y) = 1 - d(w, y) \geq 0$ . Thus  $v, y \in S$  doubly resolve  $v, w$ . Hence  $S$  doubly resolves  $K_n$  and  $\psi(K_n) = n - 1$ .  $\square$

Theorem 3 and Lemma 5 imply that every graph  $G$  satisfies

$$\beta(K_n \square G) \leq \beta(G) + \max\{n - 2, 1\}. \quad (3)$$

In certain cases, this result can be improved as follows.

**Lemma 6** *For every graph  $G$  and for all  $n \geq 1$ ,*

$$\beta(K_n \square G) \leq \max\{n - 1, 2 \cdot \beta(G)\}.$$

**Proof.** Let  $S$  be a metric basis of  $G$ . Fix a vertex  $r$  of  $K_n$ . As illustrated in Figure 1, there is a set  $T$  of  $\max\{n - 1, 2|S|\}$  vertices of  $G \square H$  such that:

- (a) for all vertices  $a \in V(K_n) \setminus \{r\}$ , there is at least one vertex  $x \in S$  for which  $ax \in T$ , and
- (b) for all  $x \in S$ , there are at least two vertices  $a, b \in V(K_n)$  for which  $ax \in T$  and  $bx \in T$ .

To prove that  $T$  resolves  $K_n \square G$ , consider two vertices  $av$  and  $bw$  of  $K_n \square G$ .




Fig. 1. The resolving set  $T$  of  $K_9 \square G$  in Lemma 6.

First suppose that  $v \neq w$ . Then there is a vertex  $x \in S$  that resolves  $v$  and  $w$  in  $G$ . Hence  $d_G(v, x) < d_G(w, x)$  without loss of generality. By (b) there are distinct vertices  $c, d \in V(K_n)$  for which  $cx \in T$  and  $dx \in T$ . If  $c \neq a$  and  $c \neq b$ , then

$$d(av, cx) = d_G(v, x) + 1 < d_G(w, x) + 1 = d(bw, cx));$$

that is,  $cx$  resolves  $av$  and  $bw$  in  $K_n \square G$ . Similarly, if  $d \neq a$  and  $d \neq b$ , then  $dx$  resolves  $av$  and  $bw$ . Otherwise  $c = a$  or  $c = b$ , and  $d = a$  or  $d = b$ . Since  $c \neq d$ , without loss of generality  $c = a$  and  $d = b$ . Then

$$d(av, cx) = d_G(v, x) < d_G(w, x) < d_G(w, x) + 1 = d(bw, cx),$$

and again  $cx$  resolves  $av$  and  $bw$  in  $K_n \square G$ .

Now suppose that  $v = w$ . Then  $a \neq b$ , and  $a \neq r$  without loss of generality. By (a) there is an  $x \in S$  for which  $ax \in T$ . Then  $d(av, ax) = d_G(v, x) < d_G(w, x) + 1 = d(bw, ax)$ ; that is,  $ax$  resolves  $av$  and  $bw$  in  $K_n \square G$ .  $\square$

When  $n$  is large in comparison with  $\beta(G)$  we know  $\beta(K_n \square G)$  exactly.

**Theorem 7** For every graph  $G$  and for all  $n \geq 2 \cdot \beta(G) + 1$ ,

$$\beta(K_n \square G) = n - 1.$$

**Proof.** The lower bound  $\beta(K_n \square G) \geq n - 1$  follows from Corollary 2 and (2). The upper bound  $\beta(K_n \square G) \leq n - 1$  is a special case of Lemma 6.  $\square$

## 6 Mastermind and Hamming Graphs

*Mastermind* is a game for two players, the *code setter* and the *code breaker*<sup>1</sup>. The code setter chooses a secret vector  $s = [s_1, s_2, \dots, s_n] \in \{1, 2, \dots, k\}^n$ . The task of the code breaker is to infer the secret vector by a series of questions, each a vector  $t = [t_1, t_2, \dots, t_n] \in \{1, 2, \dots, k\}^n$ . The code setter answers with two integers, first being the number of positions in which the secret vector and the question agree, denoted by  $a(s, t) = |\{i : s_i = t_i, 1 \leq i \leq n\}|$ . The second integer  $b(s, t)$  is the maximum of  $a(\tilde{s}, t)$ , where  $\tilde{s}$  ranges over all permutations of  $s$ .

In the commercial version of the game,  $n = 4$  and  $k = 6$ . The secret vector and each question is represented by 4 pegs each coloured with one of 6 colours. Each answer is represented by  $a(s, t)$  black pegs, and  $b(s, t) - a(s, t)$  white pegs. Knuth [22] showed that four questions suffice to determine  $s$  in this case. Here the code breaker may determine each question in response to the previous answers. *Static mastermind* is the variation in which all the questions must be supplied at once. Let  $g(n, k)$  denote the maximum, taken over all vectors  $s$ , of the minimum number of questions required to determine  $s$  in this static setting.

The *Hamming graph*  $H_{n,k}$  is the cartesian product of cliques

$$H_{n,k} = \underbrace{K_k \square K_k \square \dots \square K_k}_n.$$

Note that the hypercube  $Q_n = H_{n,2}$ . The vertices of  $H_{n,k}$  can be thought of as vectors in  $\{1, 2, \dots, k\}^n$ , with two vertices being adjacent if they differ in precisely one coordinate. Thus the distance  $d_H(v, w)$  between two vertices  $v$  and  $w$  is the number of coordinates in which their vectors differ. That is,

$$d_H(v, w) = n - a(v, w).$$

Suppose for the time being that we remove the second integer  $b(s, t)$  from the answers given by the code setter in the static mastermind game. For the code breaker to correctly infer the secret vector  $s$  from a set of questions  $T$ ,  $s$  must be uniquely determined by the values  $\{a(s, t) : t \in T\}$ . Equivalently, for any two vertices  $v$  and  $w$  of  $H_{n,k}$ , there is a  $t \in T$  for which  $a(v, t) \neq a(w, t)$ ; that is, the distances  $d_H(v, t) \neq d_H(w, t)$ . Hence the secret vector can be inferred if and only if  $T$  resolves  $H_{n,k}$ , and thus  $g(n, k) = \beta(H_{n,k})$  in this simplified

---

<sup>1</sup> Chvátal [11] referred to the code setter and code breaker as S.F. and P.G.O.M. (in honour of P.E.).



game. In this setting, Chvátal [11] proved the upper bound

$$\beta(H_{n,k}) \leq (2 + \epsilon)n \frac{1 + 2 \log k}{\log n - \log k}$$

for large  $n > n(\epsilon)$  and small  $k < n^{1-\epsilon}$ . For  $k \in \{3, 4\}$ , improvements to the constant in the above upper bound are stated without proof by Kabatianski *et al.* [20]. With the second integer  $b(s, t)$  included in the game, we have

$$g(n, k) \leq \beta(H_{n,k}).$$

Thus lower bounds for  $g(n, k)$  are also lower bounds on  $\beta(H_{n,k})$ . Kabatianski *et al.* [20] state that a ‘straightforward generalisation’ of the lower bound on  $\beta(Q_n)$  by Erdős and Rényi [13] gives for large  $n$ ,

$$\beta(H_{n,k}) \geq (2 + o(1)) \frac{n \log k}{\log n}.$$

Here we study  $\beta(H_{n,k})$  for large values of  $k$  rather than for large values of  $n$ . A similar approach is taken by Goddard [15, 16] for static Mastermind, who proved that  $g(2, k) = \lceil \frac{2}{3}k \rceil$  and  $g(3, k) = k - 1$ . Our contribution is to determine the exact value of  $\beta(H_{2,k})$ . We show that for all  $k \geq 1$ ,

$$\beta(H_{2,k}) = \left\lfloor \frac{2}{3}(2k - 1) \right\rfloor. \quad (4)$$

Equation (4) is a special case (with  $m = n = k$ ) of the following more general result.

**Theorem 8** *For all  $n \geq m \geq 1$  we have*

$$\beta(K_n \square K_m) = \begin{cases} \left\lfloor \frac{2}{3}(n + m - 1) \right\rfloor & , \text{ if } m \leq n \leq 2m - 1 \\ n - 1 & , \text{ if } n \geq 2m - 1 . \end{cases}$$

Note that two vertices of  $K_n \square K_m$  are adjacent if and only if they are in a common row or column. Otherwise they are at distance two. Fix a set  $S$  of vertices of  $K_n \square K_m$ . With respect to  $S$ , a row or column is *empty* if it contains no vertex in  $S$ , and a vertex  $v \in S$  is *lonely* if  $v$  is the only vertex of  $S$  in its row and in its column. As illustrated in Figure 2, we have the following characterisation of resolving sets in  $K_n \square K_m$ .

**Lemma 9** *For  $m, n \geq 2$ , a set  $S$  of vertices resolves  $K_n \square K_m$  if and only if:*

- (a) *there is at most one empty row and at most one empty column,*
- (b) *there is at most one lonely vertex, and*
- (c) *if there is an empty row and an empty column, then there is no lonely vertex.*




Fig. 2. Resolving set of  $K_7 \square K_7$  with one empty row, one empty column, and no lonely vertex.

**Proof.** ( $\implies$ ) First suppose that  $S$  resolves  $K_n \square K_m$ . By Corollary 2, the projections of  $S$  respectively resolve  $K_m$  and  $K_n$ . By (2), there is at most one empty row and at most one empty column. Thus (a) holds.

Suppose on the contrary that  $v$  and  $w$  are two lonely vertices in  $S$ . Thus  $v$  and  $w$  are in distinct rows and columns, and no other vertex of  $S$  is in a row or column that contains  $v$  or  $w$ . Let  $x$  be the vertex in the row of  $v$  and the column of  $w$ . Let  $y$  be the vertex in the column of  $v$  and the row of  $w$ . Then  $d(x, v) = d(y, v) = 1$ ,  $d(x, w) = d(y, w) = 1$ , and  $d(x, u) = d(y, u) = 2$  for every vertex  $u \in S \setminus \{v, w\}$ . Thus  $S$  does not resolve  $x$  and  $y$ . This contradiction proves that  $S$  satisfies (b).

Finally, suppose that there is an empty row, an empty column, and a lonely vertex  $v \in S$ . Let  $x$  be the vertex in the row of  $v$  and in the empty column. Let  $y$  be the vertex in the column of  $v$  and in the empty row. We have  $d(x, v) = d(y, v) = 1$ , and  $d(x, u) = d(y, u) = 2$  for every vertex  $u \in S \setminus \{v\}$ . Thus  $S$  does not resolve  $x$  and  $y$ . This contradiction proves that  $S$  satisfies (c).

( $\impliedby$ ) Now suppose that  $S$  is a set of vertices satisfying (a), (b) and (c). We will prove that  $S$  resolves any two vertices  $x$  and  $y$ . If  $x \in S$ , then  $x$  resolves  $x, y$ . If  $y \in S$ , then  $y$  resolves  $x, y$ . Now suppose that  $x \notin S$  and  $y \notin S$ .

If  $x$  and  $y$  are in the same row, then at least one of the columns of  $x$  and  $y$  contains a vertex  $v \in S$ . Suppose  $v$  is in the column of  $x$ . Thus  $d(x, v) = 1$  and  $d(y, v) = 2$ , and  $v$  resolves  $x, y$ . Similarly, if  $x$  and  $y$  are in the same column, then some  $v \in S$  resolves  $x, y$ .

Suppose now that  $x$  and  $y$  are in distinct rows and columns. Then there is a vertex of  $S$  in the column of  $x$  or in the column of  $y$ . Suppose  $v \in S$  is in the column of  $x$ . If  $v$  is not in the row of  $y$ ,  $d(x, v) = 1 \neq 2 = d(y, v)$ , and  $v$  resolves  $x, y$ . If  $v$  is in the row of  $y$ , by (b) and (c), at least one of the vertices in the rows and columns of  $x$  and  $y$ , but not in the intersection of two of them,

is in  $S$ . This vertex resolves  $x$  and  $y$ .  $\square$

**Lemma 10** For all  $n, m \geq 3$ ,

$$\beta(K_n \square K_m) = 2 + \min\{\beta(K_{n-2} \square K_{m-1}), \beta(K_{n-1} \square K_{m-2})\}.$$

**Proof.** We first prove that

$$\beta(K_n \square K_m) \leq 2 + \min\{\beta(K_{n-2} \square K_{m-1}), \beta(K_{n-1} \square K_{m-2})\}. \quad (5)$$

Without loss of generality  $\beta(K_{n-2} \square K_{m-1}) \leq \beta(K_{n-1} \square K_{m-2})$ . Let  $S$  be a metric basis of  $K_{n-2} \square K_{m-1}$ . Construct  $S' \subseteq V(K_n \square K_m)$  from  $S$  by adding two new vertices that are positioned in one new row and in two new columns. The number of empty rows, empty columns, and lonely vertices is the same in  $S$  and  $S'$ . Since  $S$  resolves  $K_{n-2} \square K_{m-1}$ ,  $S'$  resolves  $K_n \square K_m$  by Lemma 9. Thus  $\beta(K_n \square K_m) \leq |S'| = |S| + 2 = 2 + \beta(K_{n-2} \square K_{m-1})$ , which implies (5). It remains to prove that

$$\min\{\beta(K_{n-2} \square K_{m-1}), \beta(K_{n-1} \square K_{m-2})\} \leq \beta(K_n \square K_m) - 2. \quad (6)$$

Let  $S$  be a metric basis of  $K_n \square K_m$ . By Lemma 9,  $S$  contains two vertices  $v$  and  $w$  in the same row or column. Without loss of generality,  $v$  and  $w$  are in the same row. Construct  $S' \subseteq V(K_{n-2} \square K_{m-1})$  from  $S$  by deleting the row containing  $v$  and  $w$ , and by deleting the two columns containing  $v$  and  $w$ . The number of empty rows, empty columns, and lonely vertices is the same in  $S$  and  $S'$ . Since  $S$  resolves  $K_n \square K_m$ ,  $S'$  resolves  $K_{n-2} \square K_{m-1}$  by Lemma 9. Thus  $\beta(K_{n-2} \square K_{m-1}) \leq |S'| \leq |S| - 2 = \beta(K_n \square K_m) - 2$ , which implies (6).  $\square$

**Proof of Theorem 8.** We proceed by induction on  $n + m$  in increments of 3. (Formally speaking, we are doing induction on  $\lfloor \frac{1}{3}(n + m) \rfloor$ .)

First observe that for  $m = 1$ , we know that  $\beta(K_n \square K_m) = n - 1$ . For  $m = 2$ , we have  $\beta(K_2 \square K_2) = 2 = \lfloor \frac{2}{3}(2 + 2 - 1) \rfloor$ ,  $\beta(K_3 \square K_2) = 2 = \lfloor \frac{2}{3}(3 + 2 - 1) \rfloor$ , and  $\beta(K_n \square K_2) = n - 1$  for all  $n \geq 3$ . Thus the assertion is true for  $m \leq 2$ . Now suppose that  $m \geq 3$ . By Lemma 10 we have

$$\beta(K_n \square K_m) = 2 + \min\{\beta(K_{n-2} \square K_{m-1}), \beta(K_{n-1} \square K_{m-2})\}. \quad (7)$$

**Case 1.**  $n \geq 2m - 1$ : Then  $n \geq 2 \cdot \beta(K_m) + 1$  by (2), and  $\beta(K_n \square K_m) = n - 1$  by Theorem 7 with  $G = K_m$ .

**Case 2.**  $n = 2m - 2$ : First consider  $K_{n'} \square K_{m'}$ , where  $n' = n - 1 = 2m - 3$  and  $m' = m - 2$ . Then  $m' \leq n'$  and  $n' \geq 2m' - 1$ . By induction,

$$\beta(K_{n'} \square K_{m'}) = n' - 1 = n - 2 = \lfloor \frac{2}{3}(n + m - 1) \rfloor - 2.$$

Now consider  $K_{n'} \square K_{m'}$ , where  $m' = m - 1$  and  $n' = n - 2 = 2m - 4$ . Then  $m' \leq n' \leq 2m' - 1$ . By induction

$$\beta(K_{n'} \square K_{m'}) = \lfloor \frac{2}{3}(n' + m' - 1) \rfloor = \lfloor \frac{2}{3}(n + m - 1) \rfloor - 2.$$

By (7),  $\beta(K_n \square K_m) = \lfloor \frac{2}{3}(n + m - 1) \rfloor$ .

**Case 3.**  $n = 2m - 3$ : First consider  $K_{n'} \square K_{m'}$ , where  $m' = m - 2$  and  $n' = n - 1 = 2m - 4$ . Then  $m' \leq n'$  and  $n' \geq 2m' - 1$ . By induction,

$$\beta(K_{n'} \square K_{m'}) = n' - 1 = n - 2 = \lfloor \frac{2}{3}(n + m - 1) \rfloor - 2.$$

Now consider  $K_{n'} \square K_{m'}$ , where  $m' = m - 1$ ,  $n' = n - 2 = 2m - 5$ . For  $m \geq 4$ , we have  $m' \leq n' \leq 2m' - 1$ . By induction

$$\beta(K_{n'} \square K_{m'}) = \lfloor \frac{2}{3}(n' + m' - 1) \rfloor = \lfloor \frac{2}{3}(n + m - 1) \rfloor - 2.$$

For  $m = 4$ , we have  $n = 2m - 3 = 5$  and  $m' = m - 1 = 3$ ,  $n' = n - 2 = 3$ . We verify directly that  $\beta(K_3 \square K_3) = 3 = \lfloor \frac{2}{3}(3 + 3 - 1) \rfloor = \lfloor \frac{2}{3}(5 + 4 - 1) \rfloor - 2$ . In all cases we obtain  $\beta(K_n \square K_m) = \lfloor \frac{2}{3}(n + m - 1) \rfloor$  by (7).

**Case 4.**  $n \leq 2m - 4$ : First consider  $K_{n'} \square K_{m'}$ , where  $m' = m - 2$  and  $n' = n - 1 \leq 2m - 5$ . Then,  $m' \leq n' \leq 2m' - 1$ . By induction,

$$\beta(K_{n'} \square K_{m'}) = \lfloor \frac{2}{3}(n' + m' - 1) \rfloor = \lfloor \frac{2}{3}(n + m - 1) \rfloor - 2.$$

Now consider  $K_{n'} \square K_{m'}$ , where  $m' = m - 1$  and  $n' = n - 2 \leq 2m - 6$ . If  $m \leq n - 1$ , then  $m' \leq n' < 2m' - 1$ , and by induction

$$\beta(K_{n'} \square K_{m'}) = \lfloor \frac{2}{3}(n' + m' - 1) \rfloor = \lfloor \frac{2}{3}(n + m - 1) \rfloor - 2.$$

If  $m = n \geq 4$ , then  $n' \leq m' \leq 2n' - 1$  and by induction

$$\beta(K_{m'} \square K_{n'}) = \lfloor \frac{2}{3}(m' + n' - 1) \rfloor = \lfloor \frac{2}{3}(n + m - 1) \rfloor - 2.$$

Finally, if  $m = n = 3$ , then  $\beta(K_{n'} \square K_{m'}) = \beta(K_2 \square K_1) = 1 = \lfloor \frac{2}{3}(3 + 3 - 1) \rfloor - 2$ . In all cases, we obtain  $\beta(K_n \square K_m) = \lfloor \frac{2}{3}(m + n - 1) \rfloor$  by (7).  $\square$

## 7 Paths and Grids

Let  $P_n$  denote the path on  $n \geq 1$  vertices. Khuller *et al.* [21] and Chartrand *et al.* [8] proved that an  $n$ -vertex graph  $G$  has

$$\beta(G) = 1 \iff G = P_n. \quad (8)$$

Thus, by Theorem 7, for all  $n \geq 3$ ,

$$\beta(K_n \square P_m) = n - 1. \quad (9)$$

Minimum doubly resolving sets in paths are easily characterised.

**Lemma 11** *For all  $n \geq 2$  we have  $\psi(P_n) = 2$ . Moreover, the two endpoints of  $P_n$  are in every doubly resolving set of  $P_n$ .*

**Proof.** By definition  $\psi(G) \geq 2$  for every graph  $G \neq K_1$ . Let  $P_n = (v_1, v_2, \dots, v_n)$ . For all  $1 \leq i < j \leq n$ , we have  $d(v_i, v_1) - d(v_j, v_1) = (i - 1) - (j - 1) = i - j$ , and  $d(v_i, v_n) - d(v_j, v_n) = (n - i) - (n - j) = j - i$ . Thus  $\{v_1, v_n\}$  doubly resolve  $P_n$ , and  $\psi(P_n) = 2$ . Finally, observe that  $v_1$  is in every doubly resolving set, as otherwise  $v_1$  and  $v_2$  would not be doubly resolved. Similarly  $v_n$  is in every doubly resolving set.  $\square$

**Lemma 12** *If  $\beta(G \square H) = 2$ , then  $G$  or  $H$  is a path.*

**Proof.** Say  $S = \{av, bw\}$  resolves  $G \square H$ . Suppose that  $a = b$ . Then the projection of  $S$  onto  $G$  is a single vertex. By Lemma 1, the projection of  $S$  onto  $G$  resolves  $G$ , and by (8), only paths have singleton resolving sets. Thus  $G$  is a path, and we are done. Similarly, if  $v = w$  then  $H$  is a path, and we are done. Now suppose that  $a \neq b$  and  $v \neq w$ . Let  $c$  be the neighbour of  $b$  on a shortest path from  $a$  to  $b$ . Then  $d_G(a, c) + 1 = d_G(a, b)$  and  $d_G(b, c) = 1$ . Similarly, let  $x$  be the neighbour of  $w$  on a shortest path from  $v$  to  $w$ . Then  $d_H(v, x) + 1 = d_H(v, w)$  and  $d_H(x, w) = 1$ . This implies that  $S$  does not resolve  $bx$  and  $cw$ , since

$$d(bx, av) = d_G(a, b) + d_H(x, v) = d_G(a, c) + d_H(v, w) = d(cw, av)$$

and

$$d(bx, bw) = d_H(x, w) = 1 = d_G(b, c) = d(cw, bw).$$

This contradiction proves the result.  $\square$

Theorem 3 and Lemma 11 imply that every graph  $G$  satisfies

$$\beta(G) \leq \beta(G \square P_n) \leq \beta(G) + 1, \quad (10)$$

as proved by Chartrand *et al.* [8] in the case that  $n = 2$ .

An  $n$ -dimensional *grid* is a cartesian product of paths  $P_{m_1} \square P_{m_2} \square \cdots \square P_{m_n}$ . Equations (8) and (10) imply that,

$$\beta(P_{m_1} \square P_{m_2} \square \cdots \square P_{m_n}) \leq n. \quad (11)$$

as proved by Khuller *et al.* [21], who in addition claimed that  $\beta(P_{m_1} \square P_{m_2} \square \cdots \square P_{m_n}) = n$ . They wrote ‘we leave it for the reader to see why  $n$  is a lower bound’. This claim is false if every  $m_i = 2$  and  $n$  is large, since  $\beta(P_2 \square P_2 \square \cdots \square P_2) \rightarrow 2n/\log n$  as discussed in Section 2. Sebő and Tannier [31] claimed without proof that ‘using a result of Lindström [24]’ one can prove that

$$\limsup_{n \rightarrow \infty} \beta(P_k \square P_k \square \cdots \square P_k) \cdot \frac{\log n}{n \log k} \leq 2. \quad (12)$$

## 8 Cycles

Let  $C_n$  denote the cycle on  $n \geq 3$  vertices. Two vertices  $v$  and  $w$  of  $C_n$  are *antipodal* if  $d(v, w) = \frac{n}{2}$ . Note that no two vertices are antipodal in an odd cycle.

**Lemma 13** ([21, 30]) *For all  $n \geq 3$  we have  $\beta(C_n) = 2$ . Moreover, two vertices resolve  $C_n$  if and only if they are not antipodal.*

**Lemma 14** *For all  $n \geq 3$  we have*

$$\psi(C_n) = \begin{cases} 2 & , \text{ if } n \text{ is odd} \\ 3 & , \text{ if } n \text{ is even.} \end{cases}$$

**Proof.** We have  $\psi(C_n) \geq 2$  by definition. Now we prove the upper bound. Denote  $C_n = (v_1, v_2, \dots, v_n)$ . Consider two vertices  $v_i$  and  $v_j$  of  $C_n$ . Without loss of generality  $i < j$ .

**Case 1.**  $1 \leq i < j \leq k+1$ : Then  $d(v_i, v_1) - d(v_j, v_1) = (i-1) - (j-1) = i-j$ , and  $d(v_i, v_{k+1}) - d(v_j, v_{k+1}) = (k+1-i) - (k+1-j) = j-i \neq i-j$ . Thus  $v_1, v_{k+1}$  doubly resolve  $v_i, v_j$ .

**Case 2.**  $k+1 \leq i < j \leq n$ : Then  $d(v_i, v_1) - d(v_j, v_1) = (n+1-i) - (n+1-j) = j-i$ , and  $d(v_i, v_{k+1}) - d(v_j, v_{k+1}) = (i-k-1) - (j-k-1) = i-j \neq j-i$ . Thus  $v_1, v_{k+1}$  doubly resolve  $v_i, v_j$ .

**Case 3.**  $1 \leq i \leq k+1 < j \leq n$ : Suppose that  $v_1, v_{k+1}$  does not doubly resolve  $v_i, v_j$ . That is,  $d(v_i, v_1) - d(v_j, v_1) = d(v_i, v_{k+1}) - d(v_j, v_{k+1})$ . Thus

$(i-1) - (n+1-j) = (k+1-i) - (j-k-1)$ . Hence  $n = 2i + 2j - 2k - 4$  is even.

Therefore for odd  $n$ ,  $\{v_1, v_{k+1}\}$  doubly resolves  $C_n$ , and  $\psi(C_n) = 2$ .

For even  $n$ , in Case 3, suppose that  $v_1, v_2$  does not doubly resolve  $v_i, v_j$ . That is,  $d(v_i, v_1) - d(v_j, v_1) = d(v_i, v_2) - d(v_j, v_2)$ . Thus  $(i-1) - (n+1-j) = (i-2) - (n+2-j)$  and  $-2 = -4$ , a contradiction. Hence for even  $n$ ,  $\{v_1, v_2, v_{k+1}\}$  doubly resolve  $C_n$ , and  $\psi(C_n) \leq 3$ .

It remains to prove that  $\psi(C_n) \geq 3$  for even  $n$ . Suppose that  $\psi(C_n) \leq 2$  for some even  $n = 2k$ . By symmetry we can assume that  $\{v_1, v_i\}$  doubly resolves  $C_n$  for some  $2 \leq i \leq k+1$ .

**Case 1.**  $2 \leq i \leq k-1$ : Then  $d(v_{i+1}, v_1) - d(v_{i+2}, v_1) = i - (i+1) = -1$ , and  $d(v_{i+1}, v_i) - d(v_{i+2}, v_i) = 1 - 2 = -1$ . Thus  $v_1, v_i$  does not resolve  $v_{i+1}, v_{i+2}$ .

**Case 2.**  $i = k$ : Then  $d(v_2, v_1) - d(v_{n-1}, v_1) = 1 - 2 = -1$ , and  $d(v_2, v_i) - d(v_{n-1}, v_i) = (k-2) - (k-1) = -1$ . Thus  $v_1, v_i$  does not resolve  $v_2, v_{n-1}$ .

**Case 3.**  $i = k+1$ : Then  $d(v_2, v_1) - d(v_n, v_1) = 1 - 1 = 0$ , and  $d(v_2, v_i) - d(v_n, v_i) = (k-1) - (k-1) = 0$ . Thus  $v_1, v_i$  does not resolve  $v_2, v_n$ .

In each case we have derived a contradiction. Thus  $\psi(C_n) \geq 3$  for even  $n$ .  $\square$

Theorem 3 and Lemma 14 imply that every graph  $G$  satisfies

$$\beta(G) \leq \beta(G \square C_n) \leq \begin{cases} \beta(G) + 1 & , \text{ if } n \text{ is odd} \\ \beta(G) + 2 & , \text{ if } n \text{ is even.} \end{cases} \quad (13)$$

In particular,

$$\beta(C_m \square C_n) \leq \begin{cases} 3 & , \text{ if } m \text{ or } n \text{ is odd} \\ 4 & , \text{ otherwise.} \end{cases} \quad (14)$$

**Theorem 15** *For every graph  $G$  and for all  $n \geq 3$ , we have  $\beta(G \square C_n) = 2$  if and only if  $G$  is a path and  $n$  is odd.*

**Proof.** ( $\Leftarrow$ ) Since  $G$  is a path,  $\beta(G) = 1$  by (8). Since  $n$  is odd,  $\psi(C_n) = 2$  by Lemma 14. Thus  $\beta(G \square C_n) \leq \psi(C_n) + \beta(G) - 1 = 2$  by Theorem 3.

( $\Rightarrow$ ) Suppose that  $\beta(G \square C_n) = 2$ . Say  $S = \{av, bw\}$  resolves  $G \square C_n$ . Then  $G$  is a path by Lemma 12. It remains to show that  $n$  is odd. Suppose on the contrary that  $n = 2r$  is even. Let  $C = C_n$ . By Corollary 2, the projection  $\{v, w\}$  of  $S$  onto  $C$  resolves  $C$ . By Lemma 13, we have  $\beta(C) = 2$ , and thus  $v \neq w$ .

Moreover,  $v$  and  $w$  are not antipodal. That is,  $d_C(v, w) \leq r - 1$ . Hence there is a neighbour  $x$  of  $w$  in  $C$  with  $d_C(v, x) = d_C(v, w) + 1$ . Now consider  $G$ . If  $a \neq b$ , then using the argument from the proof of Lemma 12, we can construct a pair of vertices that are not resolved by  $S$ . So now assume  $a = b$ . That is, our resolving set is contained in a single column of  $G \square C_n$ . Let  $p$  be a neighbour of  $a$  in  $G$ . Then  $S$  does not resolve  $pw$  and  $ax$ , since  $d(pw, bw) = 1 = d(ax, bw)$  and  $d(pw, av) = 1 + d_C(v, w) = d_C(x, v) = d(ax, av)$ . This contradiction proves the result.  $\square$

By Lemma 14 and (8), we have  $\beta(P_m \square C_n) \leq \psi(C_n) + \beta(P_m) - 1 \leq 3 + 1 - 1 = 3$ . Thus Theorem 15 implies that for all  $m \geq 2$  and  $n \geq 3$  we have

$$\beta(P_m \square C_n) = \begin{cases} 2 & , \text{ if } n \text{ is odd} \\ 3 & , \text{ if } n \text{ is even.} \end{cases} \quad (15)$$

**Theorem 16** *For all  $n \geq 1$  and  $m \geq 3$  we have*

$$\beta(K_n \square C_m) = \begin{cases} 2 & , \text{ if } n = 1 \\ 2 & , \text{ if } n = 2 \text{ and } m \text{ is odd} \\ 3 & , \text{ if } n = 2 \text{ and } m \text{ is even} \\ 3 & , \text{ if } n = 3 \\ 3 & , \text{ if } n = 4 \text{ and } m \text{ is even} \\ 4 & , \text{ if } n = 4 \text{ and } m \text{ is odd} \\ n - 1 & , \text{ if } n \geq 5. \end{cases}$$

**Proof.** The case  $n \geq 2\beta(C_n) + 1 = 5$  is an immediate corollary of Theorem 7 and Lemma 13. The case  $n = 3$  is a special case of (14) since  $K_3 = C_3$ . The case  $n = 2$  is a special case of (15) since  $K_2 = P_2$ . The case  $n = 1$  is a repetition of Lemma 13.

It remains to prove the case  $n = 4$ . Say  $V(K_4) = \{a, b, c, d\}$ . First note that  $\beta(K_4 \square C_m) \geq \beta(K_4) = 3$  by Corollary 2 and (2). By Lemma 5 we have  $\psi(K_4) = 3$ . Thus  $\beta(K_4 \square C_m) \leq 4$  by Lemma 13 and Theorem 3 with  $H = K_4$ . For even  $m$ , it is easily verified that  $\{av, bv, cw\}$  resolves  $K_4 \square C_m$  for any edge  $vw$  of  $C_m$ .

It remains to prove that  $\beta(K_4 \square C_m) \geq 4$  for odd  $m = 2h + 1$ . Consider the vertices of  $K_4 \square C_m$  to be in a  $4 \times m$  grid, where two vertices in the same row are adjacent, and two vertices in the same column are adjacent if and only if they are consecutive rows or they are in the first and last rows. Suppose on the contrary that  $S = \{u, v, w\}$  resolves  $K_4 \square C_m$ . Then  $u, v, w$  are in three different columns and in at least two different rows (by considering the projections of  $S$  onto  $K_4$  and  $C_m$ ).



**Case 1.** Suppose that two vertices in  $S$ , say  $u$  and  $v$ , are in the same row. Consider the grid centred at the row of  $u, v$ . Without loss of generality,  $u$  and  $v$  are in the first and second columns, and  $w$  is in a row above  $u$  and  $v$ . Let  $x$  and  $y$  be the vertices shown in Figure 3(a). Then  $d(x, u) = d(y, u) = h + 1$ ,  $d(x, v) = d(y, v) = h + 1$ , and  $d(x, w) = d(y, w) = a$ . Thus  $S$  does not resolve  $x$  and  $y$ , which is the desired contradiction.




Fig. 3. Illustration for Theorem 16.

**Case 2.** Now suppose that  $u, v, w$  are in different rows. Without loss of generality,  $u$  is in the middle row and the first column, and  $v$  is in the second column and in a row below  $u$ , and  $w$  is in the third column and in a row above  $u$ . Let  $x$  and  $y$  be the vertices shown in Figure 3(b). Then  $d(x, u) = d(y, u) = h + 1$ ,  $d(x, v) = d(y, v) = b$ , and  $d(x, w) = d(y, w) = a$ . Thus  $S$  does not resolve  $x$  and  $y$ , which is the desired contradiction.  $\square$

## 9 Trees

Let  $v$  be a vertex of a tree  $T$ . Let  $\ell_v$  be the number of components of  $T \setminus v$  that are (possibly edgeless) paths. Khuller *et al.* [21] and Chartrand *et al.* [8] proved that for every tree  $T$  that is not a path,

$$\beta(T) = \sum_{v \in V(T)} \max\{\ell_v - 1, 0\}. \quad (16)$$

A *leaf* of a graph is a vertex of degree one. The following result for doubly resolving sets in trees is a generalisation of Lemma 11 for paths.

**Lemma 17** *The set of leaves  $L$  is the unique minimum doubly resolving set for a tree  $T$ , and  $\psi(T) = |L|$ .*

**Proof.** Every pair of vertices  $v, w$  of  $T$  lie on a path whose endpoints are leaves  $x, y$ . Clearly  $x, y$  doubly resolve  $v, w$ . Thus  $L$  is a doubly resolving set. Say  $v$  is a leaf of  $T$  whose neighbour is  $w$ . Every shortest path from  $v$  passes through  $w$ . Thus  $v, w$  can only be doubly resolved by a pair including  $v$ . Thus  $v$  is in every doubly resolving set of  $T$ . The result follows.  $\square$

Theorem 3 and Lemma 17 imply that for every tree  $T$  with  $k$  leaves and for every graph  $G$ ,

$$\beta(T \square G) \leq \beta(G) + k - 1. \quad (17)$$

Moreover, many leaves force up the metric dimension of a cartesian product.

**Lemma 18** *Every graph  $G$  with  $k \geq 2$  leaves satisfies  $\beta(G \square G) \geq k$ .*

**Proof.** Let  $S$  with a metric basis of  $G \square G$ . Let  $b$  and  $w$  be distinct leaves of  $G$  respectively adjacent to  $a$  and  $v$ . There is a vertex  $xy \in S$  that resolves  $aw$  and  $bv$ . Suppose on the contrary that  $x \neq b$  and  $y \neq w$ . Thus  $d_G(b, x) = d_G(a, x) + 1$  and  $d_G(w, y) = d_G(v, y) + 1$ . Hence  $d_G(a, x) - d_G(b, x) = d_G(v, y) - d_G(w, y) = -1$ , which implies that  $d_G(a, x) + d_G(w, y) = d_G(b, x) + d_G(v, y)$ . That is,  $d(aw, xy) = d(bv, xy)$ . Thus  $xy$  does not resolve  $aw$  and  $bv$ . This contradiction proves that  $x = b$  or  $y = w$ . Thus for every pair of leaves  $b, w$  there is a vertex  $by$  or  $xw$  in  $S$ . Suppose that for some leaf  $b$ , there is no vertex  $by \in S$ . Then for every leaf  $w$ , there is a vertex  $xw \in S$ , and  $|S| \geq k$ . Otherwise for every leaf  $b$ , there is a vertex  $by \in S$ , and again  $|S| \geq k$ .  $\square$

The following result implies that  $\psi$  is not bounded by any function of metric dimension.

**Theorem 19** *For every integer  $n \geq 3$  there is a tree  $B_n$  with  $\beta(B_n) = 2$  and*

$$n = \psi(B_n) \leq \beta(B_n \square B_n) \leq n + 1.$$

**Proof.** Let  $B_n$  be the *comb* graph obtained by attaching one leaf at every vertex of  $P_n$ . Now  $\ell_v = 1$  for every internal vertex  $v$  of  $P_n$ ;  $\ell_v = 2$  for each endpoint  $v$  of  $P_n$ ; and  $\ell_v = 0$  for every leaf  $v$  of  $P_n$ . Thus  $\beta(B_n) = 2$  by (16). Since  $B_n$  has  $n$  leaves, we have  $\psi(B_n) = n$  by Lemma 17. Moreover,  $\beta(B_n \square B_n) \geq n$  by Lemma 18. The upper bound  $\beta(B_n \square B_n) \leq n + 1$  follows from Theorem 3.  $\square$

Given that the proof of Theorem 19 is heavily dependent on the presence of leaves in  $B_n$ , it is tempting to suspect that such behaviour does not occur among more highly connected graphs. This is not the case.

**Theorem 20** *For all  $k \geq 1$  and  $n \geq 1$  there is a  $k$ -connected graph  $G_{n,k}$  for which  $\beta(G_{n,k}) \leq 2k$  and  $\beta(G_{n,k} \square G_{n,k}) \geq n$ .*

**Proof.** As illustrated in Figure 4, let  $G_{n,k}$  be the graph with vertex set  $\{v_i, w_i : 1 \leq i \leq 2kn\}$ , where every  $v_i w_i$  is an edge,  $v_i v_j$  is an edge whenever  $|i - j| \leq k$ , and  $w_i w_j$  is an edge whenever  $\lceil i/k \rceil = \lceil j/k \rceil$ . Note that  $G_{n,1} = B_{2n}$ . Clearly  $G_{n,k}$  is  $k$ -connected. It is easily seen that  $\{v_i, v_{2kn+1-i} : 1 \leq i \leq k\}$  resolves  $G_{n,k}$ . Thus  $\beta(G_{n,k}) \leq 2k$ . Say  $S$  doubly resolves  $G_{n,k}$ . On the contrary suppose that  $S \cap \{w_{\ell k+1}, w_{\ell k+2}, \dots, w_{\ell k+k}\} = \emptyset$  for some  $\ell$  with  $0 \leq \ell \leq 2n - 1$ . This implies that  $d(w_{\ell k+1}, x) = d(v_{\ell k+1}, x) + 1$  for every vertex  $x \in S$ . Hence  $S$  does not doubly resolve  $w_{\ell k+1}$  and  $v_{\ell k+1}$ . This contradiction proves that  $S \cap \{w_{\ell k+1}, w_{\ell k+2}, \dots, w_{\ell k+k}\} \neq \emptyset$  for every  $\ell$  with  $0 \leq \ell \leq 2n - 1$ . Thus  $|S| \geq 2n$ , and  $\psi(G_{n,k}) \geq 2n$ . That  $\beta(G_{n,k} \square G_{n,k}) \geq n$  follows from Lemma 4.  $\square$

We conclude that for all  $k \geq 1$ , there is no function  $f$  such that  $\beta(G \square H) \leq f(\beta(G), \beta(H))$  for all  $k$ -connected graphs  $G$  and  $H$ .




Fig. 4. The construction in Theorem 20 with  $k = 3$  and  $n = 2$ .

## References

- [1] NOGA ALON, DMITRY N. KOZLOV, AND VAN H. VU. The geometry of coin-weighing problems. In *Proc. 37th Annual Symposium on Foundations of Computer Science (FOCS '96)*, pp. 524–532. IEEE, 1996.
- [2] ZUZANA BEERLIOVA, FELIX EBERHARD, THOMAS ERLEBACH, ALEXANDER HALL, MICHAEL HOFFMANN, MATUS MIHALAK, AND L. SHANKAR RAM. Network discovery and verification. In *Proc. 31st*

- Workshop on Graph Theoretic Concepts in Computer Science (WG'05)*,  
Lecture Notes in Comput. Sci., Springer, to appear.
- [3] ALEX BOGOMOLNY AND DON GREENWELL. Cut the knot: Invitation to Mastermind, 1999. <http://www.maa.org/editorial/knot/Mastermind.html>.
  - [4] ROBERT C. BRIGHAM, GARY CHARTRAND, RONALD D. DUTTON, AND PING ZHANG. Resolving domination in graphs. *Math. Bohem.*, 128(1):25–36, 2003.
  - [5] DAVID G. CANTOR. Determining a set from the cardinalities of its intersections with other sets. *Canad. J. Math.*, 16:94–97, 1964.
  - [6] DAVID G. CANTOR AND W. H. MILLS. Determination of a subset from certain combinatorial properties. *Canad. J. Math.*, 18:42–48, 1966.
  - [7] GLENN G. CHAPPELL, JOHN GIMBEL, AND CHRIS HARTMAN. Bounds on the metric and partition dimensions of a graph, 2003. Manuscript.
  - [8] GARY CHARTRAND, LINDA EROH, MARK A. JOHNSON, AND ORTRUD R. OELLERMANN. Resolvability in graphs and the metric dimension of a graph. *Discrete Appl. Math.*, 105(1-3):99–113, 2000.
  - [9] GARY CHARTRAND, CHRISTOPHER POISSON, AND PING ZHANG. Resolvability and the upper dimension of graphs. *Comput. Math. Appl.*, 39(12):19–28, 2000.
  - [10] GARY CHARTRAND AND PING ZHANG. The theory and applications of resolvability in graphs. A survey. In *Proc. 34th Southeastern International Conf. on Combinatorics, Graph Theory and Computing*, vol. 160 of *Congr. Numer.*, pp. 47–68. 2003.
  - [11] VAŠEK CHVÁTAL. Mastermind. *Combinatorica*, 3(3-4):325–329, 1983.
  - [12] JAMES CURRIE AND ORTRUD R. OELLERMANN. The metric dimension and metric independence of a graph. *J. Combin. Math. Combin. Comput.*, 39:157–167, 2001.
  - [13] PAUL ERDŐS AND ALFRÉD RÉNYI. On two problems of information theory. *Magyar Tud. Akad. Mat. Kutató Int. Közl.*, 8:229–243, 1963.
  - [14] PETER FRANK AND ROBERT SILVERMAN. Remarks on detection problems. *Amer. Math. Monthly*, 74:171–173, 1967.
  - [15] WAYNE GODDARD. Static mastermind. *J. Combin. Math. Combin. Comput.*, 47:225–236, 2003.
  - [16] WAYNE GODDARD. Mastermind revisited. *J. Combin. Math. Combin. Comput.*, 51:215–220, 2004.
  - [17] DON L. GREENWELL. Mastermind. *J. Recr. Math.*, 30:191–192, 1999–2000.
  - [18] RICHARD K. GUY AND RICHARD J. NOWAKOWSKI. Coin-weighing problems. *Amer. Math. Monthly*, 102(2):164, 1995.
  - [19] FRANK HARARY AND ROBERT A. MELTER. On the metric dimension of a graph. *Ars Combinatoria*, 2:191–195, 1976.
  - [20] GRIGORI KABATIANSKI, V. S. LEBEDEV, AND J. THORPE. The Mastermind game and the rigidity of Hamming spaces. In *Proc. IEEE International Symposium on Information Theory (ISIT '00)*, p. 375. IEEE,

- 2000.
- [21] SAMIR KHULLER, BALAJI RAGHAVACHARI, AND AZRIEL ROSENFELD. Landmarks in graphs. *Discrete Appl. Math.*, 70(3):217–229, 1996.
  - [22] DONALD E. KNUTH. The computer as master mind. *J. Recreational Math.*, 9(1):1–6, 1976/77.
  - [23] BERNT LINDSTRÖM. On a combinatory detection problem. I. *Magyar Tud. Akad. Mat. Kutató Int. Közl.*, 9:195–207, 1964.
  - [24] BERNT LINDSTRÖM. On a combinatorial problem in number theory. *Canad. Math. Bull.*, 8:477–490, 1965.
  - [25] BERNT LINDSTRÖM. On a combinatory detection problem. II. *Studia Sci. Math. Hungar.*, 1:353–361, 1966.
  - [26] WAI HO MOW. Multiuser coding based on detecting matrices for synchronous-CDMA systems. In *Proc. Cryptography and Coding*, vol. 1355 of *Lecture Notes in Comput. Sci.*, pp. 251–257. Springer, 1997.
  - [27] NICHOLAS PIPPENGER. An information-theoretic method in combinatorial theory. *J. Combinatorial Theory Ser. A*, 23(1):99–104, 1977.
  - [28] CHRISTOPHER POISSON AND PING ZHANG. The metric dimension of unicyclic graphs. *J. Combin. Math. Combin. Comput.*, 40:17–32, 2002.
  - [29] VARAPORN SAENPHOLPHAT AND PING ZHANG. Connected resolvability of graphs. *Czechoslovak Math. J.*, 53(128)(4):827–840, 2003.
  - [30] VARAPORN SAENPHOLPHAT AND PING ZHANG. On connected resolving decompositions in graphs. *Czechoslovak Math. J.*, 54(129)(3):681–696, 2004.
  - [31] ANDRÁS SEBŐ AND ERIC TANNIER. On metric generators of graphs. *Math. Oper. Res.*, 29(2):383–393, 2004.
  - [32] B. SHANMUKHA, B. SOORYANARAYANA, AND K. S. HARINATH. Metric dimension of wheels. *Far East J. Appl. Math.*, 8(3):217–229, 2002.
  - [33] PETER J. SLATER. Leaves of trees. In *Proc. 6th Southeastern Conf. on Combinatorics, Graph Theory, and Computing*, vol. 14 of *Congressus Numerantium*, pp. 549–559. 1975.
  - [34] PETER J. SLATER. Dominating and reference sets in a graph. *J. Math. Phys. Sci.*, 22(4):445–455, 1988.
  - [35] STAFFAN SÖDERBERG AND HAROLD S. SHAPIRO. A combinatory detection problem. *Amer. Math. Monthly*, 70:1066, 1963.
  - [36] B. SOORYANARAYANA. On the metric dimension of a graph. *Indian J. Pure Appl. Math.*, 29(4):413–415, 1998.
  - [37] B. SOORYANARAYANA AND B. SHANMUKHA. A note on metric dimension. *Far East J. Appl. Math.*, 5(3):331–339, 2001.
  - [38] S. V. YUSHMANOV. Estimates for the metric dimension of a graph in terms of the diameters and the number of vertices. *Vestnik Moskov. Univ. Ser. I Mat. Mekh.*, 103:68–70, 1987.